

# Ch. 7.3, 7.4: Vectors and Complex Numbers

Johns Hopkins University

Fall 2014


# Vectors(1)

## Definition (Vector)

A vector is a quantity with magnitude and direction. The magnitude is the length of the vector and the direction is indicated by the position of the vector and arrow head on the end.

## Example

Acceleration




# Vectors(2)

## Definition

The starting point of a vector is the **initial point** and the end point is the **terminal point**.

## Example

Vector starting at  $A$  and ending at  $B$  is denoted by  $\vec{AB}$ .




## Remark (1)

We usually deal with vectors in starting position, that is the initial point is the origin, i.e.  $A = (0, 0)$ . For these vectors we only give the terminal point.

## Example

The vector  $\vec{AB}$ , with initial point  $A = (0, 0)$  and terminal point  $B = (1, 2)$  is denoted by  $\langle 1, 2 \rangle$ .




# Vectors(3)

## Definition

**Magnitude** of a vector is the length of the vector. The magnitude of the vector  $\vec{AB}$  is denoted by  $|\vec{AB}|$ .

## Question

*What is the magnitude of the vector  $\langle 1, 2 \rangle$ .*



## Remark (2)

We denote by  $\overline{AB}$  a line segment starting from  $A$  and ending at  $B$  and by  $\vec{AB}$  the vector starting from  $A$  and ending at  $B$ .

## Remark (3)

The vector  $\vec{AB}$  the vector starting at  $A$  and ending at  $B$  is a different vector from  $\vec{BA}$  the vector starting at  $B$  and ending at  $A$ .

## Remark (4)

The vector  $\vec{AA}$  the vector starting at  $A$  and ending at  $A$  is the **zero vector**.

## Question

*What's the magnitude of the zero vector?*

- (A) 0
- (B) *depends what the starting point A is.*
- (C) *we can't measure the magnitude.*
- (D) *infinite*

## Question

What can we say about the magnitude of  $\vec{AB}$  and  $\vec{BA}$ ?

(A)  $|\vec{AB}| = |\vec{BA}|$

(B)  $|\vec{AB}| = -|\vec{BA}|$

(C)  $|\vec{AB}| \geq |\vec{BA}|$

(D) *we can't always compare them.*



# Scalar Multiplication

## Definition

Let  $k$  be a real number. We refer to it as a **scalar**. Then the vector  $k\vec{A}$  is the **scalar multiplication** of the vector  $\vec{A}$  with  $k$ .

The magnitude of  $k\vec{A}$  is the magnitude of  $\vec{A}$  multiplied by  $|k|$ .


If  $k > 0$  the direction of the vector  $k\vec{A}$  is the same as the direction of  $\vec{A}$ , if  $k < 0$  the direction of the vector  $k\vec{A}$  is opposite to the direction of  $\vec{A}$ .

This is multiplying a vector by a real number - essentially we are stretching or shrinking the vector.

# Scalar Multiplication

## Example


Sketch  $k\vec{A}$ , for  $\vec{A} = \langle 1, 2 \rangle$  and  $k = 0, 2, -1$ .



# Vector Addition - Geometric

## Definition (Parallelogram Law)


Consider two vectors  $\vec{A}$  and  $\vec{B}$  (where these have the same initial point!). Then  $\vec{A} + \vec{B}$  is the vector beginning at their common initial point in the direction (and magnitude) the diagonal of the parallelogram with sides  $\vec{A}$  and  $\vec{B}$ .



# Vector Subtraction - Geometric

## Definition


To find the difference of two vectors  $\vec{A} - \vec{B}$  we compute  $\vec{A} + (-\vec{B})$ . Recall that  $-\vec{B}$  is the vector  $\vec{B}$  pointing in the opposite direction.



# Horizontal and vertical components

## Definition


We can think of every vector  $\vec{w}$  as the sum of two vectors one lying on the x-axis and the other one on the y-axis. We refer to those as the **horizontal component**  $w_x$  and the **vertical component**  $w_y$ .



# Horizontal and vertical components

## Definition


If the vector  $\vec{w}$  is in standard position with horizontal component  $\vec{w}_x$  and vertical component  $\vec{w}_y$  then we say  $\vec{w} = \langle \pm|w_x|, \pm|w_y| \rangle$ . The signs depend on the direction of the horizontal and vertical components. This is the **component form** of the vector.



# Horizontal and vertical components

## Definition

If  $\vec{w}$  is in standard position then we refer to it as the **position vector**. The angle between the positive x-axis and the position vector is called **direction angle**.




# Finding horizontal and vertical components

Consider the right triangle with sides the vector  $\vec{w}$  and its horizontal component  $\vec{w}_x$ . Let the directional angle be  $\alpha$ . Then

$$\cos \alpha =$$

$$\sin \alpha =$$



$$|\vec{w}_x| =$$

$$|\vec{w}_y| =$$



# Horizontal and vertical components


## Definition (Restatement)

If the vector  $\vec{w}$  is in standard position with horizontal component  $\vec{w}_x$  and vertical component  $\vec{w}_y$  then we say  $\vec{w} = \langle r \cos \alpha, r \sin \alpha \rangle$ . The signs depend on the direction of the horizontal and vertical components. This is the **component form** of the vector.

# Finding horizontal and vertical components

## Example

Consider the vector  $\vec{w} = \langle 1, 1 \rangle$ , with direction angle  $\alpha = \pi/4$ . Find the horizontal and vertical components and their magnitude.



## Question

*Your opinion on using slides instead of the classical lecture.*

- (A) *I think using slides has improved the quality of the lecture.*
- (B) *I like slides more - at least I can stay awake.*
- (C) *Slides are even worse than what we have been doing until now! I like the regular lecture better.*
- (D) *I am definitely a fan of the regular lecture format.*
- (E) *I hate both... so much.*

## Recall - last class

Let  $\vec{w}$  be a vector with magnitude  $|\vec{w}| = r$  and direction angle  $\alpha$ .

- 1 We can write  $\vec{w}$  as the sum of its horizontal and vertical component  $\vec{w}_x$  and  $\vec{w}_y$ .

- 2 We have a way to compute these,

$$\begin{aligned} |\vec{w}_y| &= r \sin \alpha \\ |\vec{w}_x| &= r \cos \alpha \end{aligned}$$

- 3 Also recall that  $\vec{w} = \langle r \cos \alpha, r \sin \alpha \rangle$ .

# Finding the component form given magnitude and direction

## Example

Find the vector  $\vec{w} = \langle a, b \rangle$ , with direction angle  $\alpha = 330^\circ$  and magnitude 40.

## Theorem

Let  $\vec{A} = \langle a_1, a_2 \rangle$  and  $\vec{B} = \langle b_1, b_2 \rangle$  and  $k$  is a scalar (i.e. a real number), then

- 1  $k\vec{A} = \langle ka_1, ka_2 \rangle$  (scalar product)
- 2  $\vec{A} + \vec{B} = \langle a_1 + b_1, a_2 + b_2 \rangle$
- 3  $\vec{A} - \vec{B} = \langle a_1 - b_1, a_2 - b_2 \rangle$
- 4  $\vec{A} \cdot \vec{B} = a_1 b_1 + a_2 b_2$  (dot product)

## Remark

Note that the dot product of two vectors is a number and the scalar product is a vector.

# Operations on vectors - Algebraic


## Example

Let  $\vec{w} = \langle -3, 2 \rangle$  and  $\vec{z} = \langle 1, -1 \rangle$ . Find the following (both algebraically and geometrically - where possible)

$$\vec{w} - \vec{z} =$$

$$2\vec{w} + 3\vec{z} =$$

$$\vec{w} \cdot \vec{z} =$$



# Angle between two vectors - application of dot product

## Theorem (Angle between two vectors)

If  $\vec{A}$  and  $\vec{B}$  are two non-zero vectors and  $\alpha$  is the angle between them, then

$$\cos \alpha = \frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|}.$$

## Proof.

Apply the law of cosines to the triangle with sides  $a$ ,  $b$  and  $c$ , where  $a = |\vec{A}|$ ,  $b = |\vec{B}|$  and  $c = |\vec{A} - \vec{B}|$ . □



# Angle between two vectors - application of dot product

## Example

Find the smallest possible angle between each two pairs  $\langle -5, 9 \rangle$  and  $\langle 9, 5 \rangle$ .

# Angle between two vectors - application of dot product

## Remark

If  $\vec{A}$  and  $\vec{B}$  are two non-zero vectors and  $\alpha$  is the angle between them and if  $\vec{A} \cdot \vec{B} = 0$ , then

$$\cos \alpha = \frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|} = 0.$$

Thus  $\alpha = 90^\circ$ . That is, the vectors  $\vec{A}$  and  $\vec{B}$  are **perpendicular**.

## Remark

If  $\vec{A}$  and  $\vec{B}$  are two non-zero vectors and  $\alpha$  is the angle between them and if  $\cos \alpha = \pm 1$ , then  $\alpha = 0^\circ$  or  $\alpha = 180^\circ$ . That is, the vectors  $\vec{A}$  and  $\vec{B}$  are **parallel**.

# Unit Vectors

## Definition

Vectors with length 1 are called **unit vectors**.

Two such vectors are  $\vec{i} = \langle 1, 0 \rangle$  and  $\vec{j} = \langle 0, 1 \rangle$ .

## Question

*Which of the following is a unit vector*

- (A)  $\langle 1, 1 \rangle$
- (B)  $\langle 1/2, 1/2 \rangle$
- (C)  $\langle \sqrt{2}/2, \sqrt{2}/2 \rangle$
- (D)  $\langle 2, -1 \rangle$
- (E) *none of these*

## Remark (Important Observation)

We can write every vector  $\langle a_1, a_2 \rangle$  in the following way, called a **linear combination**:


$$\langle a_1, a_2 \rangle = a_1 \langle 1, 0 \rangle + a_2 \langle 0, 1 \rangle = a_1 \vec{i} + a_2 \vec{j}.$$

## Example

Write the vector  $\langle 2, -6 \rangle$  as a linear combination of the unit vectors  $\vec{i}$  and  $\vec{j}$ .

## Definition

Complex plane, real and imaginary axis.



## Question

What is the magnitude of the vector  $\langle a, b \rangle$ ?

(A)  $\sqrt{a^2 + b^2}$

(B)  $\sqrt{a^2 - b^2}$

(C)  $a^2 + b^2$

(D)  $b - a$

(E) depends what the starting point of  $\langle a, b \rangle$  is.


# Complex Numbers - absolute value

## Definition (Absolute value of a complex number)

The absolute value of the complex number  $a + ib$  is defined by

$$|a + ib| = \sqrt{a^2 + b^2}.$$

Note that this is the distance from the center of the complex plane to the point  $(a, b)$ .





# Complex Numbers - trig form

## Definition (Trigonometric form of complex numbers)

Consider the complex number  $z = a + ib$ . Let  $r = |a + ib| = \sqrt{a^2 + b^2}$  and let  $\alpha$  be the angle between  $\langle a, b \rangle$  and the positive x-axis. Then the trigonometric form of the complex number  $z$  is

$$z = r(\cos \alpha + i \sin \alpha).$$

# Complex Numbers - trig form

Example (Write the complex number in trig form)

Write the complex number  $-2\sqrt{3} + 2i$  in trig form.

# Complex Numbers - trig form

Example (Write the complex number in standard form)

Write the complex number  $\sqrt{2}(\cos(\pi/4) + i \sin(\pi/4))$  in the form  $a + ib$ .

# Complex Numbers - trig form

## Theorem

Let  $z_1 = r_1(\cos \alpha_1 + i \sin \alpha_1)$  and  $z_2 = r_2(\cos \alpha_2 + i \sin \alpha_2)$ , then

$$z_1 z_2 = r_1 r_2 (\cos(\alpha_1 + \alpha_2) + i \sin(\alpha_1 + \alpha_2))$$

$$\frac{z_1}{z_2} = \frac{r_1}{r_2} (\cos(\alpha_1 - \alpha_2) + i \sin(\alpha_1 - \alpha_2))$$

## Proof.

Just try to compute  $z_1 z_2$  and  $\frac{z_1}{z_2}$ . □

# Complex Numbers - trig form

## Example (Product in trig form)

Use trigonometric form to find  $z_1 z_2$ , if  $z_1 = -2 + 2i\sqrt{3}$  and  $z_2 = \sqrt{3} + i$ .

## Opinion Poll (2)

### Question

*For next week what would you prefer?*

- (A) *More slides please!*
- (B) *Enough experiments, let's go back to lecture style*
- (C) *Either way is fine.*