

ARITHMETIC MACAULAYFICATION OF PROJECTIVE SCHEMES

STEVEN DALE CUTKOSKY AND HUY TÀI HÀ

ABSTRACT. In this paper, we study arithmetic Macaulayfication of projective schemes and Rees algebras of ideals. We discuss the existence of an arithmetic Macaulayfication for projective schemes. We give a simple necessary and sufficient condition for nonsingular projective varieties to possess an arithmetic Macaulayfication (Theorem 1.5). We also show that this condition is sufficient in general, but give examples to show that it is not in general necessary. We further consider Rees algebras $\mathcal{R}_\lambda(I) = R[(I_\lambda)t]$ (truncated Rees algebras) associated to a homogeneous ideal I and show that they are Cohen-Macaulay for large λ in some important cases (Theorem 2.1 and Corollary 2.2.1).

Dedicated to Wolmer Vasconcelos on the occasion of his sixty fifth birthday.

0. INTRODUCTION

In this paper, we study arithmetic Macaulayfication of projective schemes and Rees algebras of ideals.

In the first part of the paper, we discuss the problem of arithmetic Macaulayfication of projective schemes. This is a globalization of the problem of arithmetic Macaulayfication of local rings, which was first considered by Barshay in [3], and then studied extensively by many authors, such as Goto and Shimoda [12], Goto and Yamagishi [13], Brodmann [4], Schenzel [25], Lipman [23], Aberbach [1], Kurano [22], Aberbach, Huneke and Smith [2], and finally solved by Kawasaki [21]. We give a necessary and sufficient condition for a nonsingular projective scheme over a field k of characteristic 0 to have an arithmetic Macaulayfication.

Theorem 0.1. *(Theorem 1.3) Suppose X is a nonsingular projective scheme over a field k of characteristic 0. Then, X has an arithmetic Macaulayfication if and only if $H^0(X, \mathcal{O}_X) = k$ and $H^i(X, \mathcal{O}_X) = 0$ for all $i = 1, \dots, \dim X - 1$.*

2000 *Mathematics Subject Classification.* 14M05, 13A30.

Key words and phrases. Arithmetic Macaulayfication, Rees algebra, blowing up.

Steven Dale Cutkosky was partially supported by NSF.

We show that the cohomological conditions of Theorem 1.3 are sufficient conditions for an unmixed projective scheme to have an arithmetic Macaulayfication (Theorem 1.5). This result follows from the work of Kawasaki ([20], [21]). However, we show that the cohomological conditions of Theorem 1.3 are not necessary in general (Example 1.6, Example 1.7).

In the second part of this paper, we consider a natural class of Rees algebras associated to a homogeneous ideal, the *truncated Rees algebras*. This class of Rees algebras was first considered by the second author in [14] and [15] for the defining ideal of a set of points in \mathbb{P}^2 . It gave a new tool to completely answer the question on defining equations of projective embeddings of certain rational surfaces (see [14, Section 4.3]).

Our main result of this section is Theorem 2.1, from which we can conclude results such as the following.

Corollary 0.1.1. *(Corollary 2.2.1) Suppose that X is a projective Cohen-Macaulay scheme over a field k such that $H^0(X, \mathcal{O}_X) = k$, $H^i(X, \mathcal{O}_X) = 0$ for $i > 0$ and $\mathcal{I} \subset \mathcal{O}_X$ is an ideal sheaf which is locally a complete intersection. Then*

- (1) *There exists a Cohen-Macaulay standard graded k -algebra R with $a(R) < 0$ such that $X \cong \text{Proj } R$.*
- (2) *If $I \subset R$ is a homogeneous ideal such that $\tilde{I} \cong \mathcal{I}$, then there exists $\lambda_0 \geq \delta(I) + 1$ (where $\delta(I)$ is the maximum degree of a minimal set of homogeneous generators of I) such that the truncated Rees algebra*

$$\mathcal{R}_\lambda(I) = R[(I_\lambda)t]$$

is Cohen-Macaulay for all $\lambda \geq \lambda_0$.

To prove Theorem 2.1, we combine the method of [19] for studying the local cohomology of multigraded algebras with the results of [6].

Throughout this paper, let k be a field. We follow the notations of [8] and [16].

1. ARITHMETIC MACAULAYFICATION

Suppose that X is a projective scheme over a field k . We will say that X is *arithmetically Cohen-Macaulay* if there exists a Cohen-Macaulay standard graded k algebra S such that $X \cong \text{Proj } S$.

Definition. Suppose that X is a projective scheme over a field k . An *arithmetic Macaulayfication* of X is a proper birational morphism $\pi : Y \rightarrow X$ such that Y is arithmetically Cohen-Macaulay.

We shall first prove a very basic result on arithmetically Cohen-Macaulay schemes.

Lemma 1.1. (1) *Suppose that $Y = \text{Proj } S$ is an arithmetically Cohen-Macaulay scheme.*

Then, Y is a Cohen-Macaulay scheme, $H^i(Y, \mathcal{O}_Y) = 0$ for $i = 1, \dots, \dim Y - 1$, and $H^0(Y, \mathcal{O}_Y) = k$.

(2) *Suppose that $Y = \text{Proj } S$ is a Cohen-Macaulay scheme, $H^i(Y, \mathcal{O}_Y) = 0$ for $i = 1, \dots, \dim Y - 1$ and $H^0(Y, \mathcal{O}_Y) = k$. Then, there exists an integer n_0 such that for all $n \geq n_0$, the Veronese embedding of Y by $H^0(Y, \mathcal{O}_Y(n))$ is arithmetically Cohen-Macaulay.*

Proof. Let \mathfrak{m} be the maximal homogeneous ideal of S . We have isomorphisms

$$\bigoplus_{n \in \mathbb{Z}} H^i(Y, \mathcal{O}_Y(n)) \cong H_{\mathfrak{m}}^{i+1}(S), \forall i \geq 1,$$

and an exact sequence

$$0 \rightarrow H_{\mathfrak{m}}^0(S) \rightarrow S \rightarrow \bigoplus_{n \in \mathbb{Z}} H^0(Y, \mathcal{O}_Y(n)) \rightarrow H_{\mathfrak{m}}^1(S) \rightarrow 0.$$

(1) is immediate since S is Cohen-Macaulay if and only if $H_{\mathfrak{m}}^i(S) = 0$ for $i \leq d = \dim \text{Proj } S$.

To prove (2) we first observe that $H^i(Y, \mathcal{O}_Y(n)) = 0$ for $i > 0$ and $n \gg 0$ by Serre vanishing. Since Y is Cohen-Macaulay, we also have $H^i(Y, \mathcal{O}_Y(n)) = 0$ for $i < d$ and $n \ll 0$. \square

Theorem 1.2. (Hironaka [18]) *Suppose that $\pi : Y \rightarrow X$ is a birational morphism of projective nonsingular varieties over a field k of characteristic 0. Then*

$$H^i(Y, \mathcal{O}_Y) \simeq H^i(X, \mathcal{O}_X) \forall i.$$

Proof. By resolution of indeterminacy ([18]), there exists a commutative diagram of projective morphisms

$$\begin{array}{ccc} Z & \xrightarrow{f} & Y \\ & g \searrow & \downarrow \pi \\ & & X \end{array}$$

such that g is a product of blowups of nonsingular subvarieties,

$$g : Z = Z_n \xrightarrow{g_n} Z_{n-1} \xrightarrow{g_{n-1}} \dots \xrightarrow{g_2} Z_1 \xrightarrow{g_1} Z_0 = X.$$

We have ([24] or Lemma 2.1 [6])

$$R^i g_{j*} \mathcal{O}_{Z_j} = \begin{cases} 0, & i > 0 \\ \mathcal{O}_{Z_{j-1}}, & i = 0 \end{cases}$$

Thus,

$$R^i g_* \mathcal{O}_Z = \begin{cases} 0, & \text{if } i > 0 \\ \mathcal{O}_X, & \text{if } i = 0 \end{cases} \quad (1.1)$$

and

$$g^* : H^i(X, \mathcal{O}_X) \cong H^i(Z, \mathcal{O}_Z)$$

for all i . Now, by considering the commutative diagram

$$\begin{array}{ccc} H^i(Z, \mathcal{O}_Z) & \xleftarrow{f^*} & H^i(Y, \mathcal{O}_Y) \\ & g^* \searrow & \uparrow \pi^* \\ & & H^i(X, \mathcal{O}_X) \end{array}$$

we conclude that π^* is one-to-one. To show that π^* is an isomorphism we now only need to show that f^* is also one-to-one.

By resolution of indeterminacy it also gives a new diagram

$$\begin{array}{ccc} W & \xrightarrow{\gamma} & Z \\ & \beta \searrow & \downarrow f \\ & & Y \end{array}$$

where β is a product of blowups of nonsingular subvarieties, so similarly we have

$$\beta^* : H^i(Y, \mathcal{O}_Y) \cong H^i(W, \mathcal{O}_W)$$

for all i . This implies that f^* is one-to-one, and the theorem is proved. \square

Suppose that $f : Y \rightarrow X$ is a morphism of schemes, and \mathcal{F} is a sheaf of Abelian groups on Y . From the Leray spectral sequence $H^i(X, R^j f_* \mathcal{F}) \Rightarrow H^{i+j}(Y, \mathcal{F})$, we deduce the following exact sequence

$$0 \rightarrow H^1(X, f_* \mathcal{F}) \rightarrow H^1(Y, \mathcal{F}) \rightarrow H^0(X, R^1 f_* \mathcal{F}) \rightarrow H^2(X, f_* \mathcal{F}) \rightarrow H^2(Y, \mathcal{F}). \quad (1.2)$$

In the case of nonsingular varieties over a field of characteristic 0, we have a good necessary and sufficient condition for the existence of an arithmetic Macaulayfication.

Theorem 1.3. *Suppose X is a nonsingular projective scheme over a field k of characteristic 0. Then, X has an arithmetic Macaulayfication if and only if $H^0(X, \mathcal{O}_X) = k$ and $H^i(X, \mathcal{O}_X) = 0$ for all $i = 1, \dots, \dim X - 1$.*

Proof. Suppose that X is nonsingular and there exists an arithmetic Macaulayfication $f : Y = \text{Proj } S \rightarrow X$. By Lemma 1.1, we have $H^0(Y, \mathcal{O}_Y) = k$ and $H^i(Y, \mathcal{O}_Y) = 0$ for $0 < i < \dim Y$.

Let $g : Z \rightarrow Y$ be a resolution of singularities. Set $h = f \circ g$. Then, $H^i(Z, \mathcal{O}_Z) \cong H^i(X, \mathcal{O}_X)$ for all i by Theorem 1.2.

We have sequences

$$H^i(X, \mathcal{O}_X) \xrightarrow{f^*} H^i(Y, \mathcal{O}_Y) \xrightarrow{g^*} H^i(Z, \mathcal{O}_Z).$$

$h^* = g^* \circ f^*$ is an isomorphism, so we have $H^0(X, \mathcal{O}_X) = k$ and $H^i(X, \mathcal{O}_X) = 0$ for $0 < i < \dim X = \dim Y$. The necessary condition is proved.

Now suppose that $H^0(X, \mathcal{O}_X) = k$ and $H^i(X, \mathcal{O}_X) = 0$ for $i = 1, \dots, \dim X - 1$. X is a Cohen-Macaulay scheme since X is nonsingular. Now using Lemma 1.1, we can embed X as an arithmetically Cohen-Macaulay scheme Y . The sufficient condition is proved. \square

Remark 1.4. *The same proof shows that the conclusions of Theorem 1.3 hold if X has rational singularities, over a field of characteristic zero.*

From Kawasaki's work we easily deduce a very strong criterion for the existence of an arithmetic Macaulayfication over a field of arbitrary characteristic.

Theorem 1.5. *Suppose that X is an unmixed projective scheme of dimension ≥ 1 over a field k , $H^i(X, \mathcal{O}_X) = 0$ for $1 \leq i \leq \dim X - 1$ and $H^0(X, \mathcal{O}_X) = k$. Then there exists an arithmetic Macaulayfication of X .*

Proof. $X = \text{Proj } R$ where $R = \bigoplus_{i \geq 0} R_i$ is an unmixed, standard graded k -algebra. Let V be the (reduced) closed subscheme of X of non Cohen-Macaulay points, $s = \dim V$, $d = \dim X$, z_1, \dots, z_d be homogeneous elements of R satisfying the conclusions of Lemma

5.3 [20]. Since R is unmixed $s < d - 1$ (as follows from Corollary 2.4 [21]). Let $Q_i = (z_i, \dots, z_d) \subset R$ for $1 \leq i \leq s + 1$, and $I = Q_1 \cdots Q_s Q_{s+1}^{d-s-1} \subset R$.

Suppose that $\alpha \in X$ is a closed point, $y \in R_1 - \alpha$, $x_i = \frac{z_i}{y^{\deg z_i}}$ for $1 \leq i \leq d$. Let $q_i = (x_i, \dots, x_d)$, $\beta = q_1 \cdots q_s q_{s+1}^{d-s-1} \subset R_{(\alpha)}$. We have $q_i = (Q_i)_{(\alpha)}$ and $\beta = I_{(\alpha)}$. If $\beta = R_{(\alpha)}$, then $R_{(\alpha)}$ is Cohen-Macaulay and $R_{(\alpha)}[I_{(\alpha)}t]$ is Cohen-Macaulay. If $\beta \neq R_{(\alpha)}$, then there exists l such that $x_l, \dots, x_d \in \alpha_{(\alpha)}$ and $x_{l-1} \notin \alpha_{(\alpha)}$. As in the proof of Theorem 5.1 [20], x_l, \dots, x_d is a subsystem of a p-standard system of parameters for $R_{(\alpha)}$ and $R_{(\alpha)}/(x_l, \dots, x_d)R_{(\alpha)}$ is a Cohen-Macaulay ring if $l > 1$. $R_{(\alpha)}[\beta t] = R_{(\alpha)}[qt \cdots q_s q_{s+1}^{d-s-1} t]$ is Cohen-Macaulay by Corollary 4.5 [21], since $s < d - 1$ and $((0) : x_d) = (0)$ as $R_{(\alpha)}$ is unmixed.

Let \mathcal{I} be the sheafification of I , $Y = \text{Proj}(\bigoplus_{n \geq 0} \mathcal{I}^n)$, with projection $\pi : Y \rightarrow \text{Proj} R = X$. For $\alpha \in X$ a closed point, $R^i \pi_* \mathcal{O}_{Y, \alpha} = H^i(Y_\alpha, \mathcal{O}_{Y_\alpha})$ where $Y_\alpha = \text{Proj} R_{(\alpha)}[I_{(\alpha)}t] = Y \times_X \text{Spec } \mathcal{O}_{X, \alpha}$. Since $R_{(\alpha)}[I_{(\alpha)}t]$ is Cohen-Macaulay, we have $H^i(Y_\alpha, \mathcal{O}_{Y_\alpha}) = 0$ for $i > 0$ and $(\pi_* \mathcal{O}_Y)_\alpha = R_{(\alpha)} = \mathcal{O}_{X, \alpha}$ by Theorem 4.1 [23]. Thus $R^i \pi_* \mathcal{O}_Y = 0$ for $i > 0$ and $\pi_* \mathcal{O}_Y = \mathcal{O}_X$. From the Leray spectral sequence we deduce that

$$H^i(Y, \mathcal{O}_Y) = H^i(X, \pi_* \mathcal{O}_Y) = H^i(X, \mathcal{O}_X) = 0$$

for $1 \leq i \leq d - 1$ and

$$H^0(Y, \mathcal{O}_Y) = H^0(X, \pi_* \mathcal{O}_Y) = H^0(X, \mathcal{O}_X) = k.$$

It also follows from what was shown above that if $\gamma \in Y$ is a closed point, $\mathcal{O}_{Y, \gamma}$ is Cohen-Macaulay, so that Y is a Cohen-Macaulay scheme. Lemma 1.1 now implies that $Y = \text{Proj } S$ for some Cohen-Macaulay ring S . \square

Example 1.6. *The converse of Theorem 1.5 is not true, as can be seen from the following simple example. Suppose that k is an algebraically closed field, $X = \text{Proj } S$ is the cuspidal plane curve with coordinate ring $S = k[y_0, y_1, y_2]/(y_0 y_2^2 - y_1^3)$. $H^1(X, \mathcal{O}_X) \cong k$. Let $Y = X \times \mathbb{P}_k^1$. Note that Y is a Cohen-Macaulay scheme. $H^1(Y, \mathcal{O}_Y) \cong k \neq 0$, by the Künneth formula. There is a natural resolution of singularities $\mathbb{P}_k^1 \times \mathbb{P}_k^1 \rightarrow Y$, which is an arithmetic Macaulayfication, as $\mathbb{P}_k^1 \times \mathbb{P}_k^1 \cong \text{Proj } R$, with $R = k[x_0, x_1, x_2, x_3]/(x_0 x_2 - x_1 x_3)$.*

We observe that the converse of Theorem 1.5 is true for normal projective surfaces. For if X is a projective normal surface and $f : Y \rightarrow X$ is an arithmetic Macaulayfication, then

$f_*\mathcal{O}_Y = \mathcal{O}_X$, so that $k = H^0(Y, \mathcal{O}_Y) = H^0(X, \mathcal{O}_X)$, and $H^1(X, \mathcal{O}_X) = H^1(Y, \mathcal{O}_Y) = 0$ by (1.2) and Lemma 1.1.

The following example is of a normal 3-fold X such that the converse of Theorem 1.5 is false.

Example 1.7. *There exists a normal projective 3-fold B such that $H^2(B, \mathcal{O}_B) \neq 0$ and B has an arithmetic Macaulayfication.*

Proof. In section III of [5] an example is given of an m -primary ideal I in the power series ring $\mathbb{C}[[x, y, z]]$ such that $\bigoplus_{n \geq 0} I^n$ is normal but not Cohen-Macaulay. The construction there yields an example of the desired type.

Let $\beta : A \rightarrow \mathbb{P}_{\mathbb{C}}^3$ be the morphism obtained by blowing up a point in $\mathbb{P}_{\mathbb{C}}^3$, and then blowing up the 12 points which are the intersection points of a general hypersurface on the exceptional \mathbb{P}^2 with a general cubic curve C'' on the exceptional \mathbb{P}^2 . Let C' be the strict transform of C'' on A . In section III of [5], it is shown that there exists a projective morphism $\alpha : A \rightarrow B$ such that B is normal, $\alpha(C')$ is a point Q , $A - C' \rightarrow B - Q$ is an isomorphism, $\alpha_*\mathcal{O}_A \cong \mathcal{O}_B$ and $R^1\alpha_*\mathcal{O}_A \neq 0$. Since $R^1\alpha_*\mathcal{O}_A$ is supported at the single point Q , we have $H^0(B, R^1\alpha_*\mathcal{O}_A) \neq 0$. Since $H^i(A, \mathcal{O}_A) \cong H^i(\mathbb{P}^3, \mathcal{O}_{\mathbb{P}^3})$ for all i (by Theorem 1.2), we have $H^i(A, \mathcal{O}_A) = 0$ for $i = 1, 2$ and $H^0(A, \mathcal{O}_A) = \mathbb{C}$. Hence (by Lemma 1.1) $A \cong \text{Proj } S$ where S is a Cohen-Macaulay standard graded \mathbb{C} -algebra. By (1.2), we have an isomorphism $H^2(B, \mathcal{O}_B) \cong H^0(B, R^1\alpha_*\mathcal{O}_A) \neq 0$. \square

2. TRUNCATED REES ALGEBRAS

Let R be a standard graded k -algebra, $I \subseteq R$ a homogeneous ideal. The truncated Rees algebras associated to I are defined as follows.

Definition. Suppose that $I = \bigoplus_{t \geq \alpha} I_t$ is the homogeneous decomposition of I , where $\alpha = \alpha(I)$ is the minimum degree in I . For each $\lambda \geq \alpha$, we define the *truncated Rees algebra of I at degree λ* to be the Rees algebra

$$\mathcal{R}_\lambda(I) = R[(I_\lambda)t] \subseteq R[t]$$

of the ideal generated by I_λ .

Define $\delta = \delta(I)$, the maximum degree of a minimal system of homogeneous generators of I .

We will assume that $\lambda \geq \delta$. The truncated Rees algebra $\mathcal{R}_\lambda(I)$ has a bi-gradation determined by $\deg F = (d, 0)$ if $F \in R$ is homogeneous of degree d , and $\deg t = (-\lambda, 1)$, i.e.

$$\mathcal{R}_\lambda(I)_{(p,q)} = I_{p+q\lambda}^q t^q.$$

It can be seen that

$$R = \bigoplus_{n \geq 0} R_\lambda(I)_{(n,0)}$$

as a graded subring of $\mathcal{R}_\lambda(I)$, and

$$S_\lambda = \bigoplus_{n \geq 0} R_\lambda(I)_{(0,n)}$$

is another subring of $\mathcal{R}_\lambda(I)$ which we will consider. There is a natural isomorphism $S_\lambda \cong k[I_\lambda]$.

Set $X = \text{Proj } R$, $V_\lambda = \text{Proj } R_\lambda(I)$ (with respect to the above bi-grading), $\bar{V}_\lambda = \text{Proj } S_\lambda$. We have canonical projections $\pi_1 : V_\lambda \rightarrow X$ and $\pi_2 : V_\lambda \rightarrow \bar{V}_\lambda$.

V_λ can be identified with the graph of the rational map $X \dashrightarrow \bar{V}_\lambda$ induced by the natural inclusion $k[I_\lambda] \rightarrow R$, and we have an isomorphism $V_\lambda \cong \text{Proj}(\bigoplus_{n \geq 0} \mathcal{I}^n)$, the blowup of the sheafification \mathcal{I} of I (c.f. [10]). From now on we will assume that $\lambda \geq \delta + 1$. We then also have that $\bar{V}_\lambda \cong \text{Proj}(\bigoplus_{n \geq 0} \mathcal{I}^n)$ ([6, Lemma 1.1]) so that $V_\lambda \rightarrow \bar{V}_\lambda$ is an isomorphism, and we have a natural diagram of morphisms (where π_2 is an isomorphism):

$$\begin{array}{ccc} & V_\lambda & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ X & \xleftarrow{\pi} & \bar{V}_\lambda \end{array}$$

Let $\mathcal{L} = \mathcal{I}\mathcal{O}_{\bar{V}_\lambda}$. The respective gradings on R , S_λ and $R_\lambda(I)$ are related by isomorphisms

$$\mathcal{L}^q \otimes \pi^* \mathcal{O}_X(q\lambda) \cong \mathcal{O}_{\bar{V}_\lambda}(q),$$

$$\mathcal{O}_{V_\lambda}(p, q) \cong \pi_1^* \mathcal{O}_X(p) \otimes \pi_2^* \mathcal{O}_{\bar{V}_\lambda}(q). \quad (2.1)$$

For $q \in \mathbb{Z}$, let \mathcal{M}_q be the sheafification on X of the graded R -module

$$M_q = \bigoplus_{i \geq 0} R_\lambda(I)_{(i,q)}$$

so that (since $\lambda \geq \delta$)

$$\mathcal{M}_q = \begin{cases} \mathcal{O}_X & \text{if } q = 0 \\ \mathcal{I}^q(\lambda q) & q > 0 \\ 0 & q < 0 \end{cases}$$

Thus for $p \in \mathbb{Z}$,

$$\mathcal{M}_q(p) = \begin{cases} \mathcal{O}_X(p) & \text{if } q = 0 \\ \mathcal{I}^q(\lambda q + p) & q > 0 \\ 0 & q < 0 \end{cases}$$

For $p \in \mathbb{Z}$, let \mathcal{N}_p be the sheafification on \overline{V}_λ of the graded S_λ module

$$N_p = \bigoplus_{i \geq 0} R_\lambda(I)_{(p,i)}.$$

Observe that $N_0 = k[I_\lambda] = S_\lambda$,

$$\mathcal{N}_p = \begin{cases} \mathcal{O}_{\overline{V}_\lambda} & \text{if } p = 0 \\ \pi^* \mathcal{O}_X(p) & p > 0 \\ 0 & p < 0 \end{cases}$$

Thus for $q \in \mathbb{Z}$,

$$\mathcal{N}_p(q) = \begin{cases} \pi^* \mathcal{O}_X(p) \otimes \mathcal{O}_{\overline{V}_\lambda}(q) & \text{if } p \geq 0 \\ 0 & p < 0 \end{cases}$$

Our main result in this section is to show that for a certain class of standard graded k -algebras R and homogeneous ideals I , the truncated Rees algebras $\mathcal{R}_\lambda(I)$ of I are Cohen-Macaulay for large λ .

Theorem 2.1. *Suppose that R is a Cohen-Macaulay standard graded k -algebra of positive dimension $d + 1$ that has negative a -invariance, $a(R) < 0$ (Since R is Cohen-Macaulay this is equivalent to $H_{\mathfrak{m}_1}^{d+1}(R)_p = 0 \ \forall p \geq 0$, where \mathfrak{m}_1 is the maximal ideal of R). Let $I \subseteq R$ be a homogeneous ideal, and suppose that $\lambda \geq \delta(I) + 1$.*

Let \mathcal{I} be the ideal sheaf associated to I on $X = \text{Proj } R$,

$$E \cong \text{Proj } \bigoplus_{n \geq 0} \mathcal{I}^n / \mathcal{I}^{n+1}$$

be the exceptional divisor of $\pi_1 : V_\lambda \rightarrow X$, with dualizing sheaf ω_E on E . Suppose that

$$\begin{cases} \pi_{1*} \mathcal{O}_E(-\lambda m, m) = \mathcal{I}^m / \mathcal{I}^{m+1}, \forall m \geq 0, \\ R^i \pi_{1*} \mathcal{O}_E(-\lambda m, m) = 0, \forall i > 0, m \geq 0, \\ R^i \pi_{1*} \omega_E(-\lambda m, m) = 0, \forall i > 0, m \geq 2 \end{cases} \quad (2.2)$$

Then, there exists an integer λ_0 such that for all

$$\lambda \geq \lambda_0 \geq \delta + 1$$

the truncated Rees algebra $\mathcal{R}_\lambda(I)$ is Cohen-Macaulay.

To prove Theorem 2.1, we shall combine the method of [19] for studying the local cohomology of multi-graded algebras with the results of [6]. Suppose that $\lambda \geq \delta + 1$. For convenience, denote $S_{V_\lambda} = \mathcal{R}_\lambda(I)$. We need to show that S_{V_λ} is a Cohen-Macaulay ring for $\lambda \gg 0$.

Let

$$\mathfrak{m}_1 = \bigoplus_{i>0} R_\lambda(I)_{(i,0)}$$

be the irrelevant ideal of R ,

$$\mathfrak{n}_1 = \mathfrak{m}_1 R_\lambda(I) = \bigoplus_{i>0, j \geq 0} R_\lambda(I)_{(i,j)}.$$

Let

$$\mathfrak{m}_2 = \bigoplus_{j>0} R_\lambda(I)_{(0,j)}$$

be the irrelevant ideal of S_λ ,

$$\mathfrak{n}_2 = \mathfrak{m}_2 R_\lambda(I) = \bigoplus_{i \geq 0, j > 0} R_\lambda(I)_{(i,j)}.$$

Let

$$\mathfrak{m} = \bigoplus_{i+j>0} R_\lambda(I)_{(i,j)}$$

and

$$\mathfrak{n} = \bigoplus_{i,j>0} R_\lambda(I)_{(i,j)}.$$

Then $\mathfrak{n}_1 + \mathfrak{n}_2 = \mathfrak{m}$ and $\mathfrak{n}_1 \cap \mathfrak{n}_2 = \mathfrak{n}$.

S_{V_λ} is Cohen-Macaulay if and only if

$$H_{\mathfrak{m}}^i(S_{V_\lambda}) = 0, \quad \forall i = 0, \dots, d+1,$$

where $d = \dim X = \dim S_{V_\lambda} - 2$, so that from the Mayer-Vietoris sequence of cohomologies,

$$\dots \rightarrow H_{\mathfrak{m}}^i(S_{V_\lambda}) \rightarrow H_{\mathfrak{n}_1}^i(S_{V_\lambda}) \oplus H_{\mathfrak{n}_2}^i(S_{V_\lambda}) \rightarrow H_{\mathfrak{n}}^i(S_{V_\lambda}) \rightarrow H_{\mathfrak{m}}^{i+1}(S_{V_\lambda}) \rightarrow \dots$$

we see that S_{V_λ} is Cohen-Macaulay if and only if

$$\begin{cases} H_{\mathfrak{n}_1}^i(S_{V_\lambda}) \oplus H_{\mathfrak{n}_2}^i(S_{V_\lambda}) \xrightarrow{\sim} H_{\mathfrak{n}}^i(S_{V_\lambda}), \quad \forall i = 0, \dots, d \\ H_{\mathfrak{n}_1}^{d+1}(S_{V_\lambda}) \oplus H_{\mathfrak{n}_2}^{d+1}(S_{V_\lambda}) \hookrightarrow H_{\mathfrak{n}}^{d+1}(S_{V_\lambda}) \end{cases} \quad (2.3)$$

We have isomorphisms

$$H_{\mathfrak{m}_1}^i(M_q)_p \cong H_{\mathfrak{n}_1}^i(S_{V_\lambda})_{(p,q)}$$

and

$$H_{\mathfrak{m}_2}^i(N_p)_q \cong H_{\mathfrak{n}_2}^i(S_{V_\lambda})_{(p,q)}$$

for all $p, q \in \mathbb{Z}$ (c.f. Lemma 2.1 [7]).

For $p, q \in \mathbb{Z}$, we have commutative diagrams with exact rows [19, Theorem 1.4].

$$\begin{array}{ccccccccc} 0 & \rightarrow & H_n^0(S_{V_\lambda})_{(p,q)} & \rightarrow & (S_{V_\lambda})_{(p,q)} & \rightarrow & H^0(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) & \rightarrow & H_n^1(S_{V_\lambda})_{(p,q)} & \rightarrow & 0 \\ & & \uparrow & & \uparrow \wr & & \uparrow & & \uparrow & & \\ 0 & \rightarrow & H_{n_1}^0(S_{V_\lambda})_{(p,q)} & \rightarrow & (S_{V_\lambda})_{(p,q)} & \rightarrow & H^0(X, \mathcal{M}_q(p)) & \rightarrow & H_{n_1}^1(S_{V_\lambda})_{(p,q)} & \rightarrow & 0 \end{array} \quad (2.4)$$

and isomorphisms

$$H_n^i(S_{V_\lambda})_{(p,q)} \cong H^{i-1}(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) \quad (2.5)$$

and

$$H_{n_1}^i(S_{V_\lambda})_{(p,q)} \cong H^{i-1}(X, \mathcal{M}_q(p)) \quad (2.6)$$

for all $i \geq 2$.

For $p, q \in \mathbb{Z}$, we have commutative diagrams with exact rows ([19, Theorem 1.4]).

$$\begin{array}{ccccccccc} 0 & \rightarrow & H_n^0(S_{V_\lambda})_{(p,q)} & \rightarrow & (S_{V_\lambda})_{(p,q)} & \rightarrow & H^0(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) & \rightarrow & H_n^1(S_{V_\lambda})_{(p,q)} & \rightarrow & 0 \\ & & \uparrow & & \uparrow \wr & & \uparrow & & \uparrow & & \\ 0 & \rightarrow & H_{n_2}^0(S_{V_\lambda})_{(p,q)} & \rightarrow & (S_{V_\lambda})_{(p,q)} & \rightarrow & H^0(\bar{V}_\lambda, \mathcal{N}_p(q)) & \rightarrow & H_{n_1}^1(S_{V_\lambda})_{(p,q)} & \rightarrow & 0 \end{array} \quad (2.7)$$

and isomorphisms

$$H_{n_2}^i(S_{V_\lambda})_{(p,q)} \cong H^{i-1}(\bar{V}_\lambda, \mathcal{N}_p(q)) \quad (2.8)$$

for all $i \geq 2$.

By Lemma 2.1 [6]

$$R^i \pi_{1*} \mathcal{O}_{V_\lambda}(p, q) = 0 \text{ for } i > 0, q \geq 0, p \in \mathbb{Z}$$

and

$$\pi_{1*} \mathcal{O}_{V_\lambda}(p, q) \cong \mathcal{I}^q(p + q\lambda) \text{ for } q \geq 0, p \in \mathbb{Z}.$$

Thus by the Leray spectral sequence,

$$H^i(\bar{V}_\lambda, \mathcal{O}_{\bar{V}_\lambda}(q) \otimes \pi^* \mathcal{O}_X(p)) \cong H^i(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) \cong H^i(X, \mathcal{I}^q(p + q\lambda)) \quad (2.9)$$

for $i \geq 0, q \geq 0, p \in \mathbb{Z}$.

Proposition 2.2. *There exists $\lambda_0 \geq \delta + 1$ such that for $\lambda \geq \lambda_0$,*

- (1) $H^0(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) = (S_{V_\lambda})_{(p,q)}$ for $p, q \geq 0$.
- (2) $H^0(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) = 0$ for $p, q < 0$
- (3) $H^i(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) = 0$ for $i > 0, p \geq 0, q > 0$.
- (4) $H^i(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) = 0$ for $i < d, p, q < 0$

Proof. (1) follows from (2.1) and Lemma 1.3 [6].

We now prove (2). After possibly tensoring with an extension field of k , we may suppose that k is an infinite field. Suppose that $\lambda > \delta + 1$. Then $\mathcal{O}_{V_\lambda}(-r(\lambda - (\delta + 1)), r)$ is very ample for $r \geq 1$, by (2.1) and Lemma 1.1 [6]. For $r \geq 1$, there exists $\sigma_r \in H^0(V_\lambda, \mathcal{O}_{V_\lambda}(-r(\lambda - (\delta + 1)), r))$ such that (σ_r) contains no associated primes of V_λ . Thus

$$\sigma_r : \mathcal{O}_{V_\lambda}(r(\lambda - (\delta + 1)), -r) \rightarrow \mathcal{O}_{V_\lambda}$$

are inclusions. For $s > 0$, we have inclusions

$$\mathcal{O}_{V_\lambda}(r(\lambda - (\delta + 1)) - s, -r) \rightarrow \mathcal{O}_{V_\lambda}(-s, 0)$$

which induce inclusions

$$\pi_{1*}\mathcal{O}_{V_\lambda}(r(\lambda - (\delta + 1)) - s, -r) \rightarrow \mathcal{O}_X(-s)$$

and thus inclusions

$$H^0(V_\lambda, \mathcal{O}_{V_\lambda}(r(\lambda - (\delta + 1)) - s, -r)) \rightarrow H^0(X, \mathcal{O}_X(-s))$$

$H^0(X, \mathcal{O}_X(-s)) = 0$ for $s < 0$ since R is Cohen-Macaulay. Since $\lambda > \delta + 1$, we have

$$H^0(V_\lambda, \mathcal{O}_{V_\lambda}(p, q)) = 0$$

if $p, q < 0$.

(3) follows from (2.1) and Proposition 3.1 [6].

(4) follows from (2.1), Lemma 2.2 [6] and Proposition 3.2 [6]. □

We also have

$$H^i(X, \mathcal{O}_X(p)) = 0 \text{ for } i > 0, p \geq 0 \tag{2.10}$$

since $a(R) < 0$, and

$$H_{\mathfrak{m}_1}^i(R) = 0 \text{ for } i < d + 1, \tag{2.11}$$

since R is Cohen-Macaulay, so that

$$H^0(X, \mathcal{O}_X(p)) = \begin{cases} 0 & p < 0 \\ (S_{V_\lambda})_{(p,0)} & p \geq 0. \end{cases}$$

The conditions of (2.3) now follow for $\lambda \geq \lambda_0$ by (2.9), (2.1), Proposition 2.2, (2.10), (2.11) and (2.4) - (2.8). We have thus finished the proof Theorem 2.1.

Corollary 2.2.1. *Suppose that X is a projective Cohen-Macaulay scheme over a field k such that $H^0(X, \mathcal{O}_X) = k$, $H^i(X, \mathcal{O}_X) = 0$ for $i > 0$ and $\mathcal{I} \subset \mathcal{O}_X$ is an ideal sheaf which is locally a complete intersection. Then*

- (1) *There exists a Cohen-Macaulay standard graded k -algebra R with $a(R) < 0$ such that $X \cong \text{Proj } R$.*
- (2) *If $I \subset R$ is a homogeneous ideal such that $\tilde{I} \cong \mathcal{I}$, then there exists $\lambda_0 \geq \delta(I) + 1$ (where $\delta(I)$ is the maximum degree of a minimal set of homogeneous generators of I) such that the truncated Rees algebra*

$$\mathcal{R}_\lambda(I) = R[(I_\lambda)t]$$

is Cohen-Macaulay for $\lambda \geq \lambda_0$.

Proof. (1) follows from Lemma 1.1. I satisfies the condition (2.2) by (2.1) and Example 2.3 of [6], so that (2) follows from Theorem 2.1. \square

$X = \mathbb{P}_k^n$ is an especially important example satisfying the conditions on X of the corollary. Other classes of situations where the conditions in (2.2) are satisfied can be found from [6].

Remark 2.3. *Since R is Cohen-Macaulay, the hypothesis $a(R) < 0$ in Theorem 2.1 is equivalent to $H_{\mathfrak{m}_1}^{d+1}(R)_p = 0 \ \forall p \geq 0$. This is an extra condition compared to [6, Theorem 4.1].*

Remark 2.4. *Suppose that R is Cohen-Macaulay, $I \subset R$ is homogeneous, and (2.2) holds. If there exists $\lambda \geq \delta(I) + 1$ such that $\mathcal{R}_\lambda(I)$ is Cohen-Macaulay, then $a(R) < 0$.*

Proof. Let notation be as in the proof of Theorem 2.1. The Remark follows from (2.3), and the fact that for $p \geq 0$,

$$\begin{aligned} H_{\mathfrak{n}_1}^{d+1}(S_{V_\lambda})_{(p,0)} &\cong H^d(X, \mathcal{O}_X(p)), \\ H_{\mathfrak{n}_2}^{d+1}(S_{V_\lambda})_{(p,0)} &\cong H^d(\bar{V}_\lambda, \pi^* \mathcal{O}_X(p)) \\ &\cong H^d(X, \mathcal{O}_X(p)) \end{aligned}$$

and

$$H_{\mathfrak{n}}^{d+1}(S_{V_\lambda})_{(p,0)} \cong H^d(V_\lambda, \mathcal{O}_{V_\lambda}(p, 0)) \cong H^d(X, \mathcal{O}_X(p)).$$

\square

REFERENCES

- [1] Aberbach, I.M. (1996). *Arithmetic Macaulayfications using ideals of dimension one*. Illinois J. Math. **40**, 518-526.
- [2] Aberbach, I.M., Huneke, C. and Smith, K.E. (1996). *A tight closure approach to arithmetic Macaulayfication*. Illinois J. Math. **40**, 310-329.
- [3] Barshay, J. (1973). *Graded algebras of powers of ideals generated by A -sequences*. J. Algebra **25**, 90-99.
- [4] Brodmann, M. (1983). *Local cohomology of certain Rees and form rings, I*. J. Algebra **81**, 29-57.
- [5] Cutkosky, S.D. (1990). *A new characterization of rational surface singularities*. Inventiones Math. **102**, 157-177.
- [6] Cutkosky, S.D. and Herzog, J. (1997). *Cohen-Macaulay coordinate rings of blowup schemes*. Comment. Math. Helv. **72**, 605-617.
- [7] Cutkosky, S.D., Herzog, J. and Trung, N.V. (1999). *Asymptotic behaviour of the Castelnuovo-Mumford regularity*. Compositio Math. **118**, 243-261.
- [8] Eisenbud, D. (1995). *Commutative algebra with a view toward algebraic geometry*. Graduate Text 150. Springer-Verlag. New York.
- [9] Eisenbud, D. and Goto, S. (1984). *Linear free resolutions and minimal multiplicities*. J. Algebra. **88**, 89-133.
- [10] Eisenbud, D. and Harris, J. (2000). *The Geometry of Schemes*. Grad Text in Math. 197. Springer-Verlag.
- [11] Geramita, A.V., Gimigliano, A. and Pitteloud, Y. (1995). *Graded Betti numbers of some embedded rational n -folds*. Math. Ann. **301**, 363-380.
- [12] Goto, S. and Shimoda, Y. (1980). *On Rees algebras over Buchsbaum rings*. J. Math. Kyoto Univ. **20**, 691-708.
- [13] Goto, S. and Yamagishi, K. *The theory of unconditioned strong d -sequences and modules of finite local cohomology*. Preprint.
- [14] Hà, Huy Tài (2000). *Rational surfaces, from an algebraic perspective*. PhD Thesis. Queen's University, Kingston.
- [15] Hà, H. Tài. *On the Rees algebra of certain codimension two perfect ideals*. Manuscripta Mathematica. To appear.
- [16] Hartshorne, R. (1977). *Algebraic Geometry*. Grad text in Math. 52. Springer-Verlag. New York.
- [17] Herzog, J., Popescu, D. and Trung N.V. (2001). *Gröbner bases and regularity of Rees algebras*. arXiv:math.RA/0101122.
- [18] Hironaka, H. (1964). *Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II*. Annals of Math. **79**, 109-203, 205-326.
- [19] Hyry, E. (1999). *The diagonal subring and the Cohen-Macaulay property of a multigraded ring*. Trans. Amer. Math. Soc. **351**, no 6, 2213-2232.
- [20] Kawasaki, T. (2000). *On Macaulayfication of Noetherian schemes*. Trans. Amer. Math. Soc. **352**, no. 6, 2517-2552.
- [21] Kawasaki, T. (2002). *On arithmetic Macaulayfication of local rings*. Trans. Amer. Math. Soc. **354**, no. 1, 123-149.
- [22] Kurano, K. (1997). *On Macaulayfication obtained by a blow-up whose center is an equi-multiple ideal*. with an appendix by Yamagishi K. J. Algebra. **190**, 405-434.
- [23] Lipman, J. (1994). *Cohen-Macaulayness in graded algebras*. Math. Res. Letters. **1**, 149-157.
- [24] Matsumura, H. (1959). *Geometric structure of the cohomology rings in abstract algebraic geometry*. Mem. Coll. Sci. Univ. Kyoto. (A). **32**, 33-84.
- [25] Schenzel, P. (1982). *Dualisierende Komplexe in der lokalen Algebra und Buchsbaum ringe*. Lecture Notes in Math., Vol 907. Springer-Verlag.
- [26] Vidal, O.L. (1999). *On the diagonals of a Rees algebra*. PhD Thesis. Universitat de Barcelona, Barcelona.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65201, USA

E-mail address: dale@math.missouri.edu

E-mail address: tai@math.missouri.edu