

# BOX-SHAPED MATRICES AND THE DEFINING IDEAL OF CERTAIN BLOWUP SURFACES

HÀ HUY TÀI

ABSTRACT. In this paper, we generalize the notions of a matrix and its ideal of  $2 \times 2$  minors to that of a box-shaped matrix and its ideal of  $2 \times 2$  minors, and make use of these notions to study projective embeddings of certain blowup surfaces. We prove that the ideal of  $2 \times 2$  minors of a generic box-shaped matrix is a perfect prime ideal that gives the algebraic description for the Segre embedding of the product of several projective spaces. We use the notion of the ideal of  $2 \times 2$  minors of a box-shaped matrix to give an explicit description for the defining ideal of the blowup of  $\mathbb{P}^2$  along a set of  $\binom{d+1}{2}$  ( $d \in \mathbb{Z}$ ) points in generic position, embedded into projective spaces using very ample divisors which correspond to the linear systems of plane curves going through these points.

## 0. INTRODUCTION.

Ideals of minors of a matrix have been thoroughly studied over many decades. They play a significant role in the study of projective varieties. It had been a major classical problem to show that the ideal of  $t \times t$  minors of a generic matrix is a prime and perfect ideal. The proof for a general value of  $t$  was due to Eagon and Hochster from their important work in [16]. In the first part of this paper, we generalize the notions of a matrix and its ideal of  $2 \times 2$  minors to that of a *box-shaped-matrix* and its *ideal of  $2 \times 2$  minors*. Our main theorem in this section is the following theorem.

**Theorem 0.1** (Theorem 1.5). *If  $\mathcal{A}$  is a box-shaped matrix of indeterminates, then  $I_2(\mathcal{A})$  is a prime ideal in  $\mathfrak{k}[\mathcal{A}]$  (here,  $I_2(\mathcal{A})$  is the ideal of  $2 \times 2$  minors of  $\mathcal{A}$ ).*

Coupled with previous work of Grone ([13]), we also show that the ideal of  $2 \times 2$  minors of a *generic box-shaped matrix* is the defining ideal of a Segre embedding of the product of several projective space, namely  $\mathbb{P}(V_1) \times \dots \times \mathbb{P}(V_n) \hookrightarrow \mathbb{P}(V_1 \otimes \dots \otimes V_n)$ . This geometric

---

2000 *Mathematics Subject Classification*. Primary: 13C40, 14J26, Secondary: 14E25.

Current address: Institute of Mathematics, P.O. Box 631 Bò Hồ, Hà Nội 10000, Vietnam. Email: tai@hanimath.ac.vn.

realization of the ideal of  $2 \times 2$  minors of a generic box-shaped matrix enables us to study its perfection (Theorem 1.8), its Hilbert function (Proposition 1.7), and gives a Gröbner basis (Theorem 1.11).

Box-shaped matrices not only describe the Segre embedding of the product of several projective spaces, but also provide a new tool for the study of projective embeddings of certain blowup surfaces. This study is carried out in the second part of this paper. To be more precise, let  $\mathbb{X} = \{P_1, \dots, P_s\}$  be a set of  $s$  distinct points in  $\mathbb{P}^2$ , and let  $I_{\mathbb{X}} = \bigoplus_{t \geq \alpha} I_t \subseteq R = \mathfrak{k}[w_1, w_2, w_3]$  be the homogeneous decomposition of the defining ideal of  $\mathbb{X}$ , and  $\mathbb{P}^2(\mathbb{X})$  the blowup of  $\mathbb{P}^2$  centered at  $\mathbb{X}$ . The second part of this paper studies the problem of finding systems of defining equations for  $\mathbb{P}^2(\mathbb{X})$  embedded in projective spaces by very ample divisors which correspond to the linear systems of plane curves going through the points in  $\mathbb{X}$ . This problem has also been considered by several authors in the last ten years, such as [5], [6], [7], [10], [11], [18], [19] and [20].

A great deal of work has concentrated on an important special case, when  $s = \binom{d+1}{2}$  for some positive integer  $d$  and the points in  $\mathbb{X}$  are in generic position (cf. [5], [10], [11]). In this case,

$$I_{\mathbb{X}} = I_d \oplus I_{d+1} \oplus I_{d+2} \oplus \dots$$

is generated by  $I_d$  (see [9]). We also address this situation.

It is well known that, in our situation, all the linear systems  $I_t$  (for  $t \geq d+1$ ) are very ample (cf. [3], [7]), so they all give embeddings of  $\mathbb{P}^2(\mathbb{X})$  in projective spaces. If in addition, there are no  $d$  points of  $\mathbb{X}$  lying on a line, then the linear system  $I_d$  is also very ample. Under this assumption, Gimigliano studied the embedding of  $\mathbb{P}^2(\mathbb{X})$  given by the linear system  $I_d$ , which results in a White surface ([10] and [11]). White surfaces had also been studied in the classical literature ([24] and [30]). Gimigliano showed that the defining ideal of a White surface is generated by the  $3 \times 3$  minors of a  $3 \times d$  matrix of linear forms, and its defining ideal has the same Betti numbers as that of the ideal of  $3 \times 3$  minors of a  $3 \times d$  matrix of indeterminates (which was given by the Eagon-Northcott complex). The embedding of  $\mathbb{P}^2(\mathbb{X})$  given by the linear system  $I_{d+1}$  (which results in a Room surface) was then studied in detail by Geramita and Gimigliano ([5]). Geramita and Gimigliano were able to determine the resolution of the ideals defining the Room surfaces. They also proved that the defining

ideals of the embeddings of  $\mathbb{P}^2(\mathbb{X})$  given by the linear systems  $I_t$  are generated by quadrics, for all  $t \geq d + 1$ , but they were unable to write down those generators when  $t \geq d + 2$ .

In [2] and [27], a method of finding a system of defining equations for a diagonal subalgebra from that of a bigraded algebra was given. This, together with results of [22] (which gives the equations for the Rees algebra of the ideal of a set of  $\binom{d+1}{2}$  points in generic position), makes it possible, in theory, for one to write a system of defining equations for the embeddings of  $\mathbb{P}^2(\mathbb{X})$  given by the linear systems  $I_t$  for all  $t$ . However, this method has its disadvantages as pointed out in [14]. In the second part of this paper, we generalize Geramita and Gimigliano's argument on the Room surfaces and give an explicit description of the defining ideals of the embeddings of  $\mathbb{P}^2(\mathbb{X})$  given by the linear systems  $I_t$ , for all  $t \geq d + 1$ . Our main result in this section is the following theorem.

**Theorem 0.2** (Theorem 2.6). *Suppose  $t = d + n$  ( $n \geq 1$ ). Then the projective embedding of  $\mathbb{P}^2(\mathbb{X})$  given by the linear system  $I_t$  is generated by  $\binom{n+1}{2}d$  linear forms and the  $2 \times 2$  minors of a box-shaped matrix of linear forms.*

Throughout this paper,  $\mathfrak{k}$  will be our ground field. For simplicity, we assume that  $\mathfrak{k}$  is algebraically closed and of characteristic 0 (though many of our results are true for any field  $\mathfrak{k}$ ). We also let  $\mathbb{P}^2 = \mathbb{P}_{\mathfrak{k}}^2$  be the projective plane over  $\mathfrak{k}$ .

## 1. BOX-SHAPED MATRICES AND THEIR IDEALS OF $2 \times 2$ MINORS.

The techniques we use in this section are inspired by those of [26] in his study of ideals of  $2 \times 2$  minors of a matrix.

### Box-shaped matrices

Let  $S$  be a commutative ring that contains a field  $\mathfrak{k}$ . An  $n$ -dimensional array ( $n \geq 2$ )

$$\mathcal{A} = (a_{i_1 \dots i_n})_{1 \leq i_j \leq r_j, \forall j=1, \dots, n},$$

can be realized as the box

$$\mathbf{B} = \{(i_1, \dots, i_n) | 1 \leq i_j \leq r_j, \forall j\},$$

in which each integral point  $(i_1, \dots, i_n)$  is assigned the value  $a_{i_1 \dots i_n}$ .

**Definition.** An  $n$ -dimensional array  $\mathcal{A}$ , with its box-shaped realization  $\mathbf{B}$ , is called an  $n$ -dimensional box-shaped matrix of size  $r_1 \times \dots \times r_n$ .

We associate to each box-shaped matrix  $\mathcal{A}$  of elements in  $S$  a ring  $S_{\mathcal{A}} = \mathfrak{k}[\mathcal{A}]$ , the subring of  $S$  obtained by adjoining the elements of  $\mathcal{A}$  to the field  $\mathfrak{k}$ .

**Definition.** Suppose  $\mathcal{A}$  is an  $n$ -dimensional box-shaped matrix of size  $r_1 \times \dots \times r_n$  of elements in  $S$ . For each  $l = 1, 2, \dots, n$ , we call

$$a_{i_1 \dots i_l \dots i_n} a_{j_1 \dots j_l \dots j_n} - a_{i_1 \dots i_{l-1} j_l i_{l+1} \dots i_n} a_{j_1 \dots j_{l-1} i_l j_{l+1} \dots j_n} \in S_{\mathcal{A}},$$

(where  $(i_1, \dots, i_n)$  and  $(j_1, \dots, j_n)$  are any two integral points in  $\mathbf{B}$ ), a  $2 \times 2$  minor about the  $l$ -th coordinate of  $\mathcal{A}$ . A  $2 \times 2$  minor of  $\mathcal{A}$  is a  $2 \times 2$  minor about at least one of its coordinates. We let  $I_2(\mathcal{A})$  be the ideal of  $S_{\mathcal{A}}$  generated by all the  $2 \times 2$  minors of  $\mathcal{A}$ , and call it the ideal of  $2 \times 2$  minors of the box-shaped matrix  $\mathcal{A}$ .

From now on, unless stated otherwise, we focus our attention to box-shaped matrices of indeterminates. Suppose  $\mathcal{A} = (x_{i_1 \dots i_n})_{(i_1, \dots, i_n) \in \mathbf{B}}$  is an  $n$ -dimensional generic box-shaped matrix of size  $r_1 \times \dots \times r_n$  with its box-shaped realization  $\mathbf{B}$ . For each  $l = 1, \dots, n$ , let

$$\mathcal{A}_l = (x_{i_1 \dots i_n})_{(i_1, \dots, i_n) \in \mathbf{B} \text{ and } i_l < r_l},$$

and denote by  $I_2(\mathcal{A}_l)$  its ideal of  $2 \times 2$  minors (in the appropriate ring). For each  $l = 1, \dots, n$ , we also let

$$\mathbf{B}_l = \{(i_1, \dots, i_n) \in \mathbf{B} \mid i_l = r_l\},$$

and

$$I_l = \langle I_2(\mathcal{A}_l), \{x_{i_1 \dots i_n} \mid (i_1, \dots, i_n) \in \mathbf{B}_l\} \rangle.$$

Throughout this paper, to any box-shaped matrix  $\mathcal{A}$ , we always associate box-shaped matrices  $\mathcal{A}_l$ , boxes  $\mathbf{B}_l$  and all the ideals  $I_l$  defined as above. The first crucial property of box-shaped matrices of indeterminates comes in the following lemma.

**Lemma 1.1.** Suppose  $\mathcal{A} = (x_{i_1 \dots i_n})_{(i_1, \dots, i_n) \in \mathbf{B}}$  is a box-shaped matrix of indeterminates in  $S$ . Then,

(a) For any  $l \neq s \in \{1, \dots, n\}$ , we have

$$I_l \cap I_s = \langle I_2(\mathcal{A}), \{x_{i_1 \dots i_n} \mid (i_1, \dots, i_n) \in \mathbf{B}_l \cap \mathbf{B}_s\} \rangle.$$

(b) For any distinct elements  $l_1, l_2, \dots, l_t$  of  $\{1, 2, \dots, n\}$  ( $2 \leq t \leq n$ ), we have

$$\cap_{j=1}^t I_{l_j} = \langle I_2(\mathcal{A}), \{x_{i_1 \dots i_n} \mid (i_1, \dots, i_n) \in \cap_{j=1}^t \mathbf{B}_{l_j}\} \rangle.$$

*Proof.* (a) For convenience, we denote by *LHS* and *RHS* the left hand side and the right hand side of the presented equality, respectively. It is clear that  $RHS \subseteq LHS$ . We need to show the opposite direction. Let  $F \in LHS$ . Since  $F \in I_l$ , we can write  $F = F' + F''$ , where

$$F' \in I_2(\mathcal{A}_l), \text{ and } F'' = \sum_{(i_1, \dots, i_n) \in \mathbf{B}_l} F_{i_1 \dots i_n} x_{i_1 \dots i_n}.$$

It suffices to show that  $F'' \in RHS$ .  $F''$  certainly belongs to  $I_s$ . Now, for  $(i_1, \dots, i_n) \in \mathbf{B}_l$ , we write  $F_{i_1 \dots i_n}$  in the form

$$F_{i_1 \dots i_n} = \sum_{(j_1, \dots, j_n) \in \mathbf{B}_s} G_{i_1 \dots i_n, j_1 \dots j_n} x_{j_1 \dots j_n} + G_{i_1 \dots i_n},$$

where  $G_{i_1 \dots i_n}$  is independent of the indeterminates  $\{x_{j_1 \dots j_n} \mid (j_1, \dots, j_n) \in \mathbf{B}_s\}$ . Then  $F'' = G + G'$ , where

$$G = \sum_{(i_1, \dots, i_n) \in \mathbf{B}_l} G_{i_1 \dots i_n} x_{i_1 \dots i_n},$$

and

$$\begin{aligned} G' &= \sum_{(i_1, \dots, i_n) \in \mathbf{B}_l, (j_1, \dots, j_n) \in \mathbf{B}_s} G_{i_1 \dots i_n, j_1 \dots j_n} x_{i_1 \dots i_n} x_{j_1 \dots j_n} \\ &= \sum_{\substack{(i_1, \dots, i_n) \in \mathbf{B}_l, \\ (j_1, \dots, j_n) \in \mathbf{B}_s}} \left( G_{i_1 \dots i_n, j_1 \dots j_n} X_{i_1 \dots i_n, j_1 \dots j_n} + x_{i_1 \dots i_{s-1} j_s i_{s+1} \dots i_n} T_{i_1 \dots i_n, j_1 \dots j_n} \right), \end{aligned}$$

where

$$X_{i_1 \dots i_n, j_1 \dots j_n} = x_{i_1 \dots i_n} x_{j_1 \dots j_n} - x_{i_1 \dots i_{s-1} j_s i_{s+1} \dots i_n} x_{j_1 \dots j_{s-1} i_s j_{s+1} \dots j_n},$$

and  $T_{i_1 \dots i_n, j_1 \dots j_n} \in S_{\mathcal{A}}$ . Clearly,  $X_{i_1 \dots i_n, j_1 \dots j_n}$  is a  $2 \times 2$  minor about the  $s$ -th coordinate of  $\mathcal{A}$ , and since the sum is taken on  $(i_1, \dots, i_n) \in \mathbf{B}_l$  and  $(j_1, \dots, j_n) \in \mathbf{B}_s$ , the point  $(i_1, \dots, i_{s-1}, j_s, i_{s+1}, \dots, i_n)$  belongs to  $\mathbf{B}_l \cap \mathbf{B}_s$ . Thus,  $G' \in RHS$ .

It remains to show that  $G \in RHS$ . Again, we have  $G \in I_s$ , so we can write:

$$G = H + \sum_{(j_1, \dots, j_n) \in \mathbf{B}_s} H_{j_1 \dots j_n} x_{j_1 \dots j_n},$$

where  $H \in I_2(\mathcal{A}_s)$ . We may also assume that  $H$  and  $H_{j_1 \dots j_n}$ , where  $(j_1, \dots, j_n) \in \mathbf{B}_l \cap \mathbf{B}_s$ , are independent of the indeterminates

$$\{x_{j_1 \dots j_n} \mid (j_1, \dots, j_n) \in \mathbf{B}_s \setminus (\mathbf{B}_l \cap \mathbf{B}_s)\}.$$

Then

$$G - H - \sum_{(j_1, \dots, j_n) \in \mathbf{B}_l \cap \mathbf{B}_s} H_{j_1 \dots j_n} x_{j_1 \dots j_n} = \sum_{(j_1, \dots, j_n) \in \mathbf{B}_s \setminus (\mathbf{B}_l \cap \mathbf{B}_s)} H_{j_1 \dots j_n} x_{j_1 \dots j_n}.$$

The left hand side of the above equality is independent of all the indeterminates

$$\{x_{j_1 \dots j_n} \mid (j_1, \dots, j_n) \in \mathbf{B}_s \setminus (\mathbf{B}_l \cap \mathbf{B}_s)\}.$$

Thus, both sides must be zero. This implies that

$$G = H + \sum_{(j_1, \dots, j_n) \in \mathbf{B}_l \cap \mathbf{B}_s} H_{j_1 \dots j_n} x_{j_1 \dots j_n} \subseteq RHS.$$

We have proved  $LHS \subseteq RHS$ . Thus, the given equality follows.

(b) We will use induction on  $t$ . For  $t = 2$  the equality is proved in part (a). Suppose  $t > 2$ , and the equality is true for  $t - 1$ . We then have

$$\cap_{j=1}^{t-1} I_j = \langle I_2(\mathcal{A}), \{x_{i_1 \dots i_n} \mid (i_1, \dots, i_n) \in \cap_{j=1}^{t-1} \mathbf{B}_j\} \rangle.$$

It remains to prove

$$\begin{aligned} & \langle I_2(\mathcal{A}), \{x_{i_1 \dots i_n} \mid (i_1, \dots, i_n) \in \cap_{j=1}^t \mathbf{B}_j\} \rangle = \\ & = \langle I_2(\mathcal{A}), \{x_{i_1 \dots i_n} \mid (i_1, \dots, i_n) \in \cap_{j=1}^{t-1} \mathbf{B}_j\} \rangle \cap I_t. \end{aligned}$$

We can proceed in the same lines of argument as that of the proof of part (a) to show that the equality above is indeed true. Hence, the presented equality is true for all  $2 \leq t \leq n$ .  $\square$

In particular, we obtain the following corollary.

**Corollary 1.1.1.**  $\cap_{l=1}^n I_l = \langle I_2(\mathcal{A}), x_{r_1 \dots r_n} \rangle$ .

### The prime-ideal theorem

From henceforth, we shall assume that our ring  $S$  is a domain. The primeness of  $I_2(\mathcal{A})$  for a generic box-shaped matrix  $\mathcal{A}$  comes as a consequence of a series of lemmas. Note that even though the following lemmas are generalizations to higher dimension of those given in [26], most of the proofs require more arguments than what was given for their 2-dimensional statements.

**Lemma 1.2.** *Suppose  $F(\dots, x_{i_1 \dots i_n}, \dots)$  is an element of  $S_{\mathcal{A}} = \mathfrak{k}[\mathcal{A}]$ . If, for some  $x_{i_1 \dots i_n}$  of  $\mathcal{A}$ , there exists a positive integer  $\lambda$  such that  $x_{i_1 \dots i_n}^\lambda F \in I_2(\mathcal{A})$ , then, for any  $x_{j_1 \dots j_n}$  of  $\mathcal{A}$  there exists a non-negative integer  $\nu$  such that  $x_{j_1 \dots j_n}^\nu F \in I_2(\mathcal{A})$ .*

*Proof.* Denote by  $Z$  the multiplicatively closed subset of  $S_{\mathcal{A}}$  consisting of all non-negative powers of  $x_{j_1 \dots j_n}$ , and let  $S_Z$  be the localization of  $S_{\mathcal{A}}$  at the set  $Z$ . Let  $\phi : S_{\mathcal{A}} \rightarrow S_Z$  be the ring homomorphism defined by  $\phi(c) = c$  for all  $c \in \mathfrak{k}$ , and

$$\phi(x_{i_1 \dots i_n}) = \frac{x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n}}{x_{j_1 \dots j_n}^{n-1}} \text{ for all } x_{i_1 \dots i_n} \in \mathcal{A}.$$

Obviously,  $\phi$  is a well-defined map. It is easy to verify that  $\phi(a) = 0$  for any  $2 \times 2$  minors  $a$  of  $\mathcal{A}$ . Thus,  $\phi(I_2(\mathcal{A})) = 0$ . Moreover,

$$x_{i_1 \dots i_n}^\lambda F(\dots, x_{i_1 \dots i_n}, \dots) \in I_2(\mathcal{A}).$$

Therefore, in  $S_Z$ ,

$$\left( \frac{x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n}}{x_{j_1 \dots j_n}^{n-1}} \right)^\lambda F(\dots, \phi(x_{i_1 \dots i_n}), \dots) = 0.$$

Since  $S_{\mathcal{A}}$  is a domain, so is  $S_Z$ . Hence,

$$F(\dots, \phi(x_{i_1 \dots i_n}), \dots) = 0 \text{ in } S_Z.$$

Now, using binomial expansions, we can write

$$F(\dots, x_{i_1 \dots i_n}, \dots) = F\left(\dots, \frac{x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n}}{x_{j_1 \dots j_n}^{n-1}}, \dots\right) + K,$$

where  $K$  belongs to the ideal of  $S_Z$  generated by elements of the form

$$x_{i_1 \dots i_n} - \frac{x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n}}{x_{j_1 \dots j_n}^{n-1}}.$$

The generators of this  $S_Z$ -ideal can be rewritten as

$$x_{i_1 \dots i_n} x_{j_1 \dots j_n}^{n-1} - x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n}.$$

We shall prove that  $K$  belongs to the ideal of  $S_Z$  generated by  $I_2(\mathcal{A})$ , or equivalently, we prove that these generators, considered as elements of  $S_{\mathcal{A}}$ , belong to  $I_2(\mathcal{A})$ . Indeed, using induction on  $n$ , modulo  $I_2(\mathcal{A})$ , we have

$$\begin{aligned} K_n &= x_{i_1 \dots i_n} x_{j_1 \dots j_n}^{n-1} - x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n} \\ &= x_{i_1 j_2 \dots j_n} x_{j_1 i_2 \dots i_n} x_{j_1 \dots j_n}^{n-2} + (x_{i_1 \dots i_n} x_{j_1 \dots j_n} - x_{i_1 j_2 \dots j_n} x_{j_1 i_2 \dots i_n}) x_{j_1 \dots j_n}^{n-2} \\ &\quad - x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n} \\ &\equiv x_{i_1 j_2 \dots j_n} x_{j_1 i_2 \dots i_n} x_{j_1 \dots j_n}^{n-2} - x_{i_1 j_2 \dots j_n} x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n} \\ &= x_{i_1 j_2 \dots j_n} K_{n-1}, \end{aligned}$$

where

$$K_{n-1} = x_{j_1 i_2 \dots i_n} x_{j_1 \dots j_n}^{n-2} - x_{j_1 i_2 j_3 \dots j_n} \cdots x_{j_1 \dots j_{n-1} i_n}.$$

Since every indeterminate appearing in the expression  $K_{n-1}$  has  $j_1$  in its first index, we can view  $K_{n-1}$  as just the same expression as  $K_n$  but given by the  $(n-1)$ -dimensional box-shaped matrix  $\mathcal{A}' = (x_{i_1 \dots i_n})_{i_1=j_1}$ . By induction hypothesis,  $K_{n-1}$  then belongs to  $I_2(\mathcal{A}') \subseteq I_2(\mathcal{A})$ . And hence,  $K_n \in I_2(\mathcal{A})$ .

We have just proved that  $K$  belongs to the ideal of  $S_Z$  generated by  $I_2(\mathcal{A})$ . Equivalently,  $F(\dots, x_{i_1 \dots i_n}, \dots)$  belongs to the ideal of  $S_Z$  generated by  $I_2(\mathcal{A})$ . Therefore, there exists a  $\nu$  such that

$$x_{j_1 \dots j_n}^\nu F(\dots, x_{i_1 \dots i_n}, \dots) \in I_2(\mathcal{A}) \text{ in } S_{\mathcal{A}}.$$

The lemma is proved. □

**Lemma 1.3.** *Suppose  $l \in \{1, 2, \dots, n\}$ . Suppose also that  $F \in S_{\mathcal{A}} = \mathfrak{k}[\mathcal{A}]$  is a polynomial independent of the indeterminates  $x_{i_1 \dots i_n}$  for all  $(i_1, \dots, i_n) \in \mathbf{B}_l$  such that  $I_2(\mathcal{A}_l) : F = I_2(\mathcal{A}_l)$ . Then  $I_l : F = I_l$ .*

*Proof.* The proof follows in exactly the same line as that of [26]. □

**Lemma 1.4.** *Let  $F \in S_{\mathcal{A}} = \mathfrak{k}[\mathcal{A}]$  and suppose that  $x_{1\dots 1}^\lambda F \in I_2(\mathcal{A})$  for some positive integer  $\lambda$ . Then  $F \in I_2(\mathcal{A})$ . In other words,  $I_2(\mathcal{A}) : x_{1\dots 1}^\lambda = I_2(\mathcal{A})$ .*

*Proof.* We use induction on  $n$ . When  $n = 2$ , the result follows from that of [26]. Suppose  $n > 2$ ,  $\mathcal{A}$  is an  $n$ -dimensional box-shaped matrix of indeterminates, and the lemma is true for any box-shaped matrices of lower dimension. We now use induction on  $r_1 + \dots + r_n$ . We may assume that  $r_i \geq 2$  for all  $i = 1, \dots, n$  (since otherwise,  $\mathcal{A}$  collapses to an  $(n - 1)$ -dimensional box-shaped matrix, and the result follows from the induction hypothesis), and the lemma is true for any  $n$ -dimensional box-shaped matrix with smaller value of  $r_1 + \dots + r_n$ .

If  $F$  is of degree zero, then  $x_{1\dots 1}^\lambda F$  belongs to the ideal of  $2 \times 2$  minors of a box-shaped matrix obtained from  $\mathcal{A}$  by letting all the indeterminates  $x_{i_1 \dots i_n}$ , for  $(i_1, \dots, i_n) \neq (1, \dots, 1)$ , be zero. Yet, this ideal is zero, so  $F = 0 \in I_2(\mathcal{A})$ . We may use induction again, assuming that the degree of  $F$  is bigger than zero, and the lemma holds for polynomials whose degrees are smaller than that of  $F$ .

Now,  $x_{1\dots 1}^\lambda F \in I_2(\mathcal{A}) \subseteq \bigcap_{j=1}^n I_j$  by Corollary 1.1.1, so in particular,  $x_{1\dots 1}^\lambda F \in I_j$  for all  $j$ . Moreover, by the induction hypothesis, we have  $I_2(\mathcal{A}_j) : x_{1\dots 1}^\lambda = I_2(\mathcal{A}_j)$ . Thus, by Lemma 1.3,  $I_j : x_{1\dots 1}^\lambda = I_j$ . This implies that  $F \in \bigcap_{j=1}^n I_j = \langle I_2(\mathcal{A}), x_{r_1 \dots r_n} \rangle$ . Write  $F = F_1 + x_{r_1 \dots r_n} F_2$ , where  $F_1 \in I_2(\mathcal{A})$ . Since  $I_2(\mathcal{A})$  is homogeneous, we may assume that the degree of  $F_2$  is smaller than that of  $F$ . We have  $x_{1\dots 1}^\lambda F = x_{1\dots 1}^\lambda F_1 + x_{1\dots 1}^\lambda x_{r_1 \dots r_n} F_2 \in I_2(\mathcal{A})$ . Thus,  $x_{r_1 \dots r_n} x_{1\dots 1}^\lambda F_2 \in I_2(\mathcal{A})$ . By Lemma 1.2, there is a non-negative integer  $\nu$  such that  $x_{1\dots 1}^{\lambda+\nu} F_2 = x_{1\dots 1}^\nu x_{1\dots 1}^\lambda F_2 \in I_2(\mathcal{A})$ . By our induction hypothesis on the degree of  $F$ , we have  $F_2 \in I_2(\mathcal{A})$ . Hence,  $F \in I_2(\mathcal{A})$  as required.  $\square$

The primeness of the ideal of  $2 \times 2$  minors of a box-shaped matrix in the generic case is stated as follows.

**Theorem 1.5.** *If  $\mathcal{A}$  is a box-shaped matrix of indeterminates, then  $I_2(\mathcal{A})$  is a prime ideal in  $\mathfrak{k}[\mathcal{A}]$ .*

*Proof.* Suppose that  $F(\dots, x_{i_1 \dots i_n}, \dots)G(\dots, x_{i_1 \dots i_n}, \dots) \in I_2(\mathcal{A})$ , where  $F, G \in S_{\mathcal{A}} = \mathfrak{k}[\mathcal{A}]$ . Let  $Z$  be the multiplicatively closed subset of  $S_{\mathcal{A}}$  consisting of all non-negative powers of

$x_{1\dots 1}$ , and let  $S_Z$  be the localization of  $S_{\mathcal{A}}$  at  $Z$ . Similar to what was done in Lemma 1.2, we define a map

$$\varphi : S_{\mathcal{A}} \rightarrow S_Z,$$

by sending  $\mathfrak{k}$  to  $\mathfrak{k}$ , and sending  $x_{i_1\dots i_n}$  to  $\frac{x_{i_1 1\dots 1} x_{1 i_2 1\dots 1} \cdots x_{1\dots 1 i_n}}{x_{1\dots 1}^{n-1}}$  for all  $x_{i_1\dots i_n} \in \mathcal{A}$ . It is easy to verify that  $\varphi(a) = 0$  for any  $2 \times 2$  minors  $a$  of  $\mathcal{A}$ . Thus,  $\varphi(I_2(\mathcal{A})) = 0$ . Moreover,  $F(\dots, x_{i_1\dots i_n}, \dots)G(\dots, x_{i_1\dots i_n}, \dots) \in I_2(\mathcal{A})$ . Hence, in  $S_Z$ ,

$$F(\dots, \varphi(x_{i_1\dots i_n}), \dots) G(\dots, \varphi(x_{i_1\dots i_n}), \dots) = 0.$$

Since  $S_{\mathcal{A}}$  is a domain, so is  $S_Z$ . Thus, at least one of the two factors has to be zero. Suppose

$$F(\dots, \frac{x_{i_1 1\dots 1} x_{1 i_2 1\dots 1} \cdots x_{1\dots 1 i_n}}{x_{1\dots 1}^{n-1}}, \dots) = 0.$$

Now, similar to what was done in Lemma 1.2, we deduce that there exists a  $\nu$  such that  $x_{1\dots 1}^\nu F(\dots, x_{i_1\dots i_n}, \dots) \in I_2(\mathcal{A})$  in  $S_{\mathcal{A}}$ . Hence, by Lemma 1.4,  $F \in I_2(\mathcal{A})$ , and this completes the proof.  $\square$

### Segre embedding, Cohen-Macaulayness and Koszul property

Suppose  $V_1, V_2, \dots, V_n$  are vector spaces of dimensions  $r_1, r_2, \dots, r_n$ , respectively. Recall the following definition.

**Definition.** A tensor  $z \in V_1 \otimes \dots \otimes V_n$  is referred to as *decomposable* if there exist  $v_j \in V_j$  for all  $j = 1, \dots, n$ , such that  $z = v_1 \otimes \dots \otimes v_n$ .

Now, let  $\{e_{j1}, \dots, e_{jr_j}\}$  be a basis for  $V_j$  for all  $j = 1, \dots, n$ . Then a basis of  $V_1 \otimes \dots \otimes V_n$  is given by

$$\{\epsilon_{i_1\dots i_n} = e_{1i_1} \otimes \dots \otimes e_{ni_n} \mid 1 \leq i_j \leq r_j \ \forall j = 1, \dots, n\}.$$

A tensor  $z \in V_1 \otimes \dots \otimes V_n$  is represented by

$$z = \sum_{i_1\dots i_n} y_{i_1\dots i_n} \epsilon_{i_1\dots i_n},$$

and a vector  $v_j \in V_j$  is given by

$$v_j = \sum_{k=1}^{r_j} u_{jk} e_{jk}.$$

Thus, to have  $z = v_1 \otimes \dots \otimes v_n$ , is the same as to have

$$y_{i_1 \dots i_n} = u_{1i_1} \dots u_{ni_n}, \text{ for all } i_1 \dots i_n.$$

This is clearly the equations describing the image of the following Segre embedding:

$$\mathbb{P}(V_1) \times \dots \times \mathbb{P}(V_n) \hookrightarrow \mathbb{P}(V_1 \otimes \dots \otimes V_n).$$

Hence, a tensor  $z \in V_1 \otimes \dots \otimes V_n$  is decomposable if and only if its corresponding point in  $\mathbb{P}(V_1 \otimes \dots \otimes V_n)$  is in the image of the above Segre embedding.

The geometric realization of the ideal of  $2 \times 2$  minors of a generic matrix  $\mathcal{A}$  comes from the work of Grone ([13]), which we rephrase in the following proposition.

**Proposition 1.6** (Grone, 1977). *Suppose  $\mathcal{A}$  is a generic box-shaped matrix of size  $r_1 \times \dots \times r_n$ , and  $V_1, \dots, V_n$  are vector spaces of dimension  $r_1, \dots, r_n$ , respectively. Then  $I_2(\mathcal{A})$  gives a set of equations that describe the decomposable tensors in  $V_1 \otimes \dots \otimes V_n$ .*

Since the Segre embedding of the product of several projective spaces is a closed immersion, Grone's result gives an immediate corollary, which demonstrates the geometric realization of  $I_2(\mathcal{A})$ .

**Corollary 1.6.1.** *If  $\mathcal{A}$  is an  $n$  dimensional generic box-shaped matrix of size  $r_1 \times \dots \times r_n$ , then  $I_2(\mathcal{A})$  gives the defining ideal of the Segre embedding*

$$\mathbb{P}(V_1) \times \dots \times \mathbb{P}(V_n) \hookrightarrow \mathbb{P}(V_1 \otimes \dots \otimes V_n).$$

where  $V_1, \dots, V_n$  are vector spaces of dimensions  $r_1, \dots, r_n$ , respectively.

*Proof.* The result follows from the fact that  $I_2(\mathcal{A})$  is a prime ideal. □

From this, we can calculate the Hilbert function of the ideal of  $2 \times 2$  minors of a generic box-shaped matrix as follows.

**Proposition 1.7.** *The Hilbert function of  $I_2(\mathcal{A})$  is*

$$\mathbf{H}(I_2(\mathcal{A}), t) = \binom{\prod_{i=1}^n r_i + t - 1}{t} - \prod_{i=1}^n \binom{r_i + t - 1}{t} \quad \forall t \geq 0.$$

*Proof.* It is easy to see that all homogeneous polynomials of degree  $t$  on  $\mathbb{P}^{\prod r_i - 1}$  restricted to the image of  $\mathbb{P}^{r_1 - 1} \times \dots \times \mathbb{P}^{r_n - 1}$  gives all multi-homogeneous polynomials of degree  $(t, \dots, t)$  in  $\mathbb{P}^{r_1 - 1} \times \dots \times \mathbb{P}^{r_n - 1}$ . Thus the Hilbert function of the homogeneous coordinate ring of the Segre embedding is  $\prod_{i=1}^n \binom{r_i + t - 1}{t}$ . The proposition now follows.  $\square$

**Remark:** It is clear that any Segre embedding is Hilbertian, i.e. its Hilbert function and its Hilbert polynomial are the same.

The geometric realization of  $I_2(\mathcal{A})$  and Propostion 1.7 give us the perfection of  $I_2(\mathcal{A})$ . The result is stated as follows.

**Theorem 1.8.** *If  $\mathcal{A}$  is an  $n$ -dimensional generic box-shaped matrix of size  $r_1 \times \dots \times r_n$ , then  $I_2(\mathcal{A})$  is a perfect ideal of grade  $\prod_{i=1}^n r_i - \sum_{i=1}^n r_i + (n - 1)$ .*

*Proof.* We let  $S_i = \mathfrak{k}[y_{i,1}, \dots, y_{i,r_i}]$  be the homogeneous coordinate ring of  $\mathbb{P}^{r_i - 1}$  for all  $i$ . Clearly,  $S_i$  is Cohen-Macaulay for all  $i$ . By results of [28, page 378] and Proposition 1.7, it follows by induction on  $n$  that the Segre product  $\otimes_{i=1}^n S_i$  is a Cohen-Macaulay ring. Furthermore, this ring is exactly the coordinate ring of the Segre embedding  $\mathbb{P}^{r_1 - 1} \times \dots \times \mathbb{P}^{r_n - 1} \hookrightarrow \mathbb{P}^{\prod r_i - 1}$ . Thus, since  $I_2(\mathcal{A})$  is the defining ideal of this Segre embedding, i.e.  $\otimes_{i=1}^n S_i \simeq \mathfrak{k}[\mathcal{A}]/I_2(\mathcal{A})$ , we have  $I_2(\mathcal{A})$  is a perfect ideal. The grade of  $I_2(\mathcal{A})$  comes from the codimension of the Segre embedding, which is exactly  $\prod_{i=1}^n r_i - \sum_{i=1}^n r_i + (n - 1)$ . The theorem is proved.  $\square$

**Remark:** The perfection of  $I_2(\mathcal{A})$  also comes from a more general result of Hochster ([17, Theorem 1]).

We have an immediate corollary.

**Corollary 1.8.1.** *Suppose  $V_1, \dots, V_n$  are vector spaces of dimensions  $r_1, \dots, r_n$ . Then, the homogeneous coordinate ring of the Segre embedding*

$$\mathbb{P}(V_1) \times \dots \times \mathbb{P}(V_n) \hookrightarrow \mathbb{P}(V_1 \otimes \dots \otimes V_n)$$

*is always Cohen-Macaulay.*

We now recall the following folklore result (cf. [21]).

**Lemma 1.9** ([21], Lemma 6.3A). *Let  $I$  be a proper ideal of  $\mathbb{Z}[x_1, \dots, x_n]$  such that  $\mathbb{Z}[x_1, \dots, x_n]/I$  is  $\mathbb{Z}$ -flat, and  $I$  is perfect of grade  $g$ . Suppose  $S$  is a Noetherian ring and  $a_1, \dots, a_n$  are elements of  $S$ . Let  $I'$  be the ideal given by the image of  $I$  under the ring homomorphism of  $\mathbb{Z}[x_1, \dots, x_n] \rightarrow S$  sending  $x_i$  to  $a_i$ . Then  $\text{grade } I' \leq g$ , and if the equality is attained then  $I'$  is a perfect ideal.*

We note also that our calculations and arguments, so far, are independent of the field  $\mathfrak{k}$ . In fact, the same calculations and arguments would apply if we have any commutative Noetherian ring with identity instead of  $\mathfrak{k}$ . Thus, our results hold when we substitute  $\mathfrak{k}$  by  $\mathbb{Z}$ , the ring of integers. This, together with Lemma 1.9, gives rise to the following result for any box-shaped matrix  $\mathcal{A}$ .

**Theorem 1.10.** *Suppose  $\mathcal{A}$  is any  $n$ -dimensional box-shaped matrix of size  $r_1 \times \dots \times r_n$ . Then,*

$$\text{grade } I_2(\mathcal{A}) \leq \prod_{i=1}^n r_i - \sum_{i=1}^n r_i + (n - 1),$$

*and if the equality is attained then  $I_2(\mathcal{A})$  is a perfect ideal.*

*Proof.* The result follows from Lemma 1.9 and the fact that our Theorem 1.8 is still true if instead of  $\mathfrak{k}$  we have the ring  $\mathbb{Z}$ . □

We return to the generic situation. Suppose again that

$$\mathcal{A} = (x_{i_1 \dots i_n})_{(i_1, \dots, i_n) \in \mathbf{B}}$$

is a generic box-shaped matrix of size  $r_1 \times \dots \times r_n$ . The following theorem gives a Gröbner basis for  $I_2(\mathcal{A})$ .

**Theorem 1.11.** *Under the degree reverse lexicographic monomial ordering on  $S_{\mathcal{A}} = \mathfrak{k}[\mathcal{A}]$ , in which the variables  $x_{i_1 \dots i_n}$  are ordered by lexicographic ordering on their indices (assuming that  $1 < 2 < \dots < n$ ), the  $2 \times 2$  minors of  $\mathcal{A}$  form a Gröbner basis for  $I_2(\mathcal{A})$ .*

*Proof.* Let  $\leq_{\text{lex}}$  be the lexicographic ordering on  $\mathbb{N}^n$ . We order the variables of  $S_{\mathcal{A}}$  by

$$x_{i_1 \dots i_n} \leq x_{j_1 \dots j_n} \Leftrightarrow (i_1, \dots, i_n) \leq_{\text{lex}} (j_1, \dots, j_n),$$

and use degree reverse lexicographic ordering on the monomials of  $S_{\mathcal{A}}$ . We shall prove that under this monomial ordering, the  $2 \times 2$  minors of  $\mathcal{A}$  form a Gröbner basis for  $I_2(\mathcal{A})$ .

Let  $\mathcal{G}$  be the collection of all  $2 \times 2$  minors of  $\mathcal{A}$ . It suffices to show that the leading terms of  $\mathcal{G}$  generate the leading term ideal of  $I_2(\mathcal{A})$ . By contradiction, suppose  $F \in I_2(\mathcal{A})$ , and  $T$ , the leading term of  $F$ , is not generated by the leading terms of  $\mathcal{G}$ . Clearly, from the nature of  $I_2(\mathcal{A})$ ,  $T$  is a monomial with at least 2 different indeterminates. We consider a new partial ordering on the indeterminates of  $S_{\mathcal{A}}$ , defined by

$$x_{i_1 \dots i_n} \preceq x_{j_1 \dots j_n} \Leftrightarrow i_l \leq j_l \quad \forall l = 1, \dots, n.$$

Suppose  $x_{i_1 \dots i_n}$  and  $x_{j_1 \dots j_n}$  are any two different indeterminates present in  $T$ . Without loss of generality, assume that  $x_{i_1 \dots i_n} < x_{j_1 \dots j_n}$ , i.e. there exists a positive integer  $u$  such that  $i_l = j_l$  for all  $l = 1, \dots, u-1$ , and  $i_u < j_u$ . It is easy to see that if  $x_{i_1 \dots i_n} \not\preceq x_{j_1 \dots j_n}$  then there exists another integer  $v > u$  such that  $i_v > j_v$ . In this case,

$$x_{i_1 \dots i_{v-1} j_v i_{v+1} \dots i_n} < x_{i_1 \dots i_n}, x_{j_1 \dots j_n} < x_{j_1 \dots j_{v-1} i_v j_{v+1} \dots j_n}.$$

Thus,  $x_{i_1 \dots i_n} x_{j_1 \dots j_n}$  is the leading term of

$$x_{i_1 \dots i_n} x_{j_1 \dots j_n} - x_{i_1 \dots i_{v-1} j_v i_{v+1} \dots i_n} x_{j_1 \dots j_{v-1} i_v j_{v+1} \dots j_n} \in \mathcal{G},$$

whence  $T$  is generated by the leading terms of  $\mathcal{G}$ , a contradiction. Hence, these two indeterminates must be comparable, i.e.  $x_{i_1 \dots i_n} \preceq x_{j_1 \dots j_n}$ . This is true for any two different indeterminates of  $T$ . Therefore,  $T$  can be rewritten as

$$T = x_{t_{11} \dots t_{1n}} x_{t_{21} \dots t_{2n}} \cdots x_{t_{p1} \dots t_{pn}},$$

for some positive integer  $p \geq 2$ , where

$$x_{t_{11} \dots t_{1n}} \preceq x_{t_{21} \dots t_{2n}} \preceq \cdots \preceq x_{t_{p1} \dots t_{pn}}.$$

Now, let  $[y_{i,1} : \dots : y_{i,r_i}]$  represent the homogeneous coordinates of  $\mathbb{P}^{r_i-1}$  for all  $i = 1, \dots, n$ . Since  $I_2(\mathcal{A})$  is the defining ideal of the Segre embedding

$$\mathbb{P}^{r_1-1} \times \cdots \times \mathbb{P}^{r_n-1} \hookrightarrow \mathbb{P}^{\prod r_i-1},$$

$F$  vanishes when we substitute the indeterminate  $x_{i_1 \dots i_n}$  by  $\prod_{l=1}^n y_{l,i_l}$  for all  $(i_1, \dots, i_n)$ . It is also clear that after this substitution,  $F$  becomes a polynomial on the variables  $y_{i,j}$ . This polynomial is zero for all values of the variables  $y_{i,j}$ , so it must be the zero polynomial (since

the ground field  $\mathfrak{k}$  is infinite). This implies that there must be a term  $T'$  of  $F$  ( $T' \neq T$ ) which cancels  $T$  after the substitution. Suppose  $x_{k_1 \dots k_n}$  is an indeterminate present in  $T'$ . Since  $T'$  cancels  $T$  after the substitution, for each  $l = 1, \dots, n$ ,  $k_l \in \{t_{1l}, \dots, t_{pl}\}$ . From the partial ordering on the indeterminates in  $T$ , it is now clear that  $k_l \geq t_{1l}$  for all  $l = 1, \dots, n$ , whence  $x_{t_{11} \dots t_{1n}} \leq x_{k_1 \dots k_n}$ . If  $x_{t_{11} \dots t_{1n}} < x_{k_1 \dots k_n}$  for every indeterminate  $x_{k_1 \dots k_n}$  in  $T'$ , then  $T < T'$ , which is a contradiction since  $T$  is the leading term of  $F$ . Otherwise, suppose  $x_{t_{11} \dots t_{1n}}$  is contained in  $T'$ , then by considering  $T/x_{t_{11} \dots t_{1n}}$  and  $T'/x_{t_{11} \dots t_{1n}}$ , and continuing the process, we eventually would, again, get a contradiction.

The theorem is proved. □

**Remark:** From the proof above, it is easy to see that the  $2 \times 2$  minors of  $\mathcal{A}$  form a Gröbner basis for  $I_2(\mathcal{A})$  under any monomial ordering on  $S_{\mathcal{A}}$  that satisfies the condition that if  $g = x_{i_1 \dots i_n} x_{j_1 \dots j_n} - x_{p_1 \dots p_n} x_{q_1 \dots q_n}$  is an element of  $\mathcal{G}$ , where  $x_{p_1 \dots p_n} \preceq x_{q_1 \dots q_n}$ , then  $x_{i_1 \dots i_n} x_{j_1 \dots j_n}$  is the leading term of  $g$ . Degree reverse lexicographic monomial ordering is merely one of those monomial orderings that satisfies this condition. We choose this ordering since it is practical in most computational algebra packages, such as CoCoA and Macaulay2.

The theorem gives rise to an interesting corollary.

**Corollary 1.11.1.** *Suppose  $V_1, \dots, V_n$  are vector spaces of dimensions  $r_1, \dots, r_n$ . Then, the homogeneous coordinate ring of the Segre embedding*

$$\mathbb{P}(V_1) \times \dots \times \mathbb{P}(V_n) \hookrightarrow \mathbb{P}(V_1 \otimes \dots \otimes V_n)$$

*is a Koszul algebra.*

*Proof.* This follows from the fact that all  $2 \times 2$  minors of  $\mathcal{A}$  are quadratic forms. □

### 3-dimensional box-shaped matrices

In the last part of this section, we briefly look at a particular class of box-shaped matrices, those of dimension 3. Besides the usual matrices, 3-dimensional box-shaped matrices are the easiest that can be visualized. To visualize all the  $2 \times 2$  minors of a 3-dimensional box-shaped matrix, one only needs to take any two lines parallel to one of the axes, and looks at their intersection with any two planes parallel to the other two axes of our fixed system of

coordinates. 3-dimensional box-shaped matrices not only describe the Segre embedding of the product of 3 projective spaces, but also give a tool in studying certain blowup surfaces, as it will be discussed in the next section. We first extend the notion of a box-shaped matrix of indeterminates to that of a *weak box-shaped matrix of indeterminates*.

**Definition.** Suppose  $\mathcal{A} = (a_{ijk})_{(i,j,k) \in \mathbf{B}}$  is a box-shaped matrix of forms in a ring  $S$ . For each integer  $l$  let  $\mathcal{A}_{(x,l)}$  be the matrix given by the collection  $\{a_{ijk} | (i,j,k) \in \mathbf{B}, i = l\}$ . We call  $\mathcal{A}_{(x,l)}$  an *x-section* of the box-shaped matrix  $\mathcal{A}$ . The y-sections and z-sections of  $\mathcal{A}$  are defined similarly.

**Definition.** A box-shaped matrix  $\mathcal{A} = (x_{ijk})_{(i,j,k) \in \mathbf{B}}$ , with box-shaped realization  $\mathbf{B}$ , of forms in a domain  $S$  is called a *weak box-shaped matrix of indeterminates* if

- (a) All the entries in  $\mathcal{A}$  are indeterminates of  $S$ , i.e. algebraically independent over  $\mathfrak{k}$ .
- (b)  $\langle I_2(\mathcal{A}), x_{r_1 r_2 r_3} \rangle = \bigcap_{l=1}^3 I_l$  where the ideals  $I_l$  are defined as that of a general  $n$ -dimensional box-shaped matrix.
- (c) There exists an integral point  $(i, j, k) \in B$  such that when we set all indeterminates other than  $x_{ijk}$  of  $\mathfrak{k}[\mathcal{A}]$  to zero, the ideal  $I_2(\mathcal{A})$  is the zero ideal.
- (d) The ideals of  $2 \times 2$  minors of sections  $\mathcal{A}_{x,i}$ ,  $\mathcal{A}_{y,j}$  and  $\mathcal{A}_{z,k}$  are prime ideals.

With this bigger class of box-shaped matrices, the primeness of their ideals of  $2 \times 2$  minors still holds.

**Proposition 1.12.**  $I_2(\mathcal{A})$  is a prime ideal in  $\mathfrak{k}[\mathcal{A}]$  for any weak box-shaped matrix of indeterminates  $\mathcal{A}$ .

*Proof.* First, we can always re-arrange the indices such that  $(i, j, k)$  becomes  $(1, 1, 1)$ . The proof now follows in the same lines as that of Theorem 1.5.  $\square$

## 2. PROJECTIVE EMBEDDINGS OF BLOWUP SURFACES.

Let  $\mathbb{X} \subseteq \mathbb{P}^2$  be a set of  $s = \binom{d+1}{2}$  points ( $d \in \mathbb{Z}, d \geq 1$ ) that are in generic position. Let  $\mathbb{P}^2(\mathbb{X})$  be the blowup of  $\mathbb{P}^2$  along the points of  $\mathbb{X}$ , and let  $I_{\mathbb{X}} = \bigoplus_{t \geq d} I_t \subseteq R = \mathfrak{k}[w_1, w_2, w_3]$  be the defining ideal of  $\mathbb{X}$ . Let  $\overline{\Lambda}_t$  be the surface obtained by embedding  $\mathbb{P}^2(\mathbb{X})$  using the

linear system  $I_t$  ( $t = d + n$ ,  $n \geq 1$ ). In this section, we give an explicit description for a system of defining equations for  $\overline{\Lambda}_t$  for any  $t$ . We start by a simple result, which could be of folklore.

**Lemma 2.1.** *Suppose  $S, R$  and  $T$  are Noetherian commutative rings with identity, and  $\phi : S \rightarrow R$  and  $\psi : R \rightarrow T$  are surjective ring homomorphisms. Suppose also that  $f_1, f_2, \dots, f_n$  are generators for  $\ker \phi \subseteq S$  and  $g_1, g_2, \dots, g_m$  are generators for  $\ker \psi \subseteq R$ . Let  $p_j$  be a preimage of  $g_j$  for all  $j$ , then  $f_1, \dots, f_n, p_1, \dots, p_m$  give a set of generators for  $\ker (\psi \circ \phi) \subseteq S$ .*

*Proof.* Clearly  $f_i$ 's and  $p_j$ 's are all in  $\ker (\psi \circ \phi)$ . Moreover, if  $x \in \ker (\psi \circ \phi)$  then either  $\phi(x) = 0$  or  $\phi(x)$  is a linear combination of the  $g_j$ 's. The result is now trivial.  $\square$

To proceed, it follows from [9] that  $I_{\mathbb{X}}$  is minimally generated in degree  $d$ . By the Hilbert-Burch theorem, these generators are the  $d \times d$  minors of a  $d \times (d + 1)$  matrix, say  $\mathbf{L}$ , of linear forms :

$$\mathbf{L} = (L_{ij}), \quad L_{ij} \in R_1 \text{ for } i = 1, 2, \dots, d \text{ and } j = 1, 2, \dots, d + 1.$$

In this notation,

$$I_{\mathbb{X}} = (F_1, \dots, F_{d+1}), \quad F_i = (-1)^{i+1} \det(\mathbf{L} \setminus i^{\text{th}} \text{ column}).$$

For  $\alpha = (\alpha_1, \alpha_2, \alpha_3)$ , we write  $w^\alpha$  for  $w_1^{\alpha_1} w_2^{\alpha_2} w_3^{\alpha_3}$ , and denote  $|\alpha| = \alpha_1 + \alpha_2 + \alpha_3$ . A system of generators of the vector space  $I_t$  is given by  $\binom{n+2}{2}(d+1)$  forms  $w^\alpha F_j$  for  $j = 1, 2, \dots, d+1$  and  $|\alpha| = n$ .

Consider the rational map

$$\varphi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^p, \quad p = \binom{n+2}{2}(d+1) - 1,$$

given by  $\varphi(P) = [w^\alpha F_j]$  (we order the  $\alpha$ 's by lexicographic ordering with  $w_1 > w_2 > w_3$ ).  $\overline{\Lambda}_t$  embedded in  $\mathbb{P}^p$  is given by the closure of the image of  $\varphi$ .

Let  $z_1 = w_1^n, z_2 = w_1^{n-1} w_2, \dots, z_u = w_3^n$ , where  $u = \binom{n+2}{2}$  (again, we arrange the terms in lexicographic order). We use homogeneous coordinates  $[x_{ij}]_{1 \leq i \leq u, 1 \leq j \leq d+1}$  of  $\mathbb{P}^p$  such that

$$\varphi([w_1 : w_2 : w_3]) = [x_{ij}], \quad \text{where } x_{ij} = z_i F_j. \quad (2.1)$$

The vector space dimension of  $I_t$  is  $(n+1)d + \binom{n+2}{2}$ , so there must be  $\binom{n+1}{2}d$  dependence relations among the  $w^\alpha F_j$ 's. Those relations can be found as follows.

Let  $\beta = (\beta_1, \beta_2, \beta_3)$  with  $|\beta| = n-1$ . For each  $l = 1, 2, \dots, d$ , we have

$$0 = \det \begin{pmatrix} w^\beta L_{l1} & w^\beta L_{l2} & \dots & w^\beta L_{ld+1} \\ & & & \mathbf{L} \end{pmatrix} = \sum_{j=1}^{d+1} L_{lj} w^\beta F_j.$$

Since  $L_{lj} = \sum_{k=1}^3 \lambda_{ljk} w_k$ , so by grouping similar terms, we get

$$\sum_{|\alpha|=n, 1 \leq j \leq d+1} \mu_{l\alpha j} w^\alpha F_j = 0, \quad \forall l = 1, 2, \dots, d,$$

where

$$\mu_{l\alpha j} = \sum_{w^\beta w_k = w^\alpha} \lambda_{ljk} \text{ for each } l, \alpha \text{ and } j.$$

These are the dependence relations of the  $w^\alpha F_j$ 's. In terms of  $z_i$ 's, we can rewrite them as

$$\sum_{i,j} \mu_{lij} z_i F_j = 0, \quad \forall l = 1, 2, \dots, d.$$

These give rise to the following equations:

$$\sum_{1 \leq i \leq u, 1 \leq j \leq d+1} \mu_{lij} x_{ij} = 0, \quad \forall l = 1, 2, \dots, d. \quad (2.2)$$

There are  $d$  relations of the form (2.2) for each  $\beta$ , and the number of such  $\beta$ 's is  $\binom{n+1}{2}$ . By abuse of notation, we denote the collection of all these  $\binom{n+1}{2}d$  relations by (2.2). The relations in (2.2) would be independent relations if we can show that the  $\binom{n+1}{2}d \times \binom{n+2}{2}(d+1)$  matrix  $\mathbf{E}$  of the coefficients  $\mu_{lij}$  has maximal rank. Indeed, we shall use a similar argument to that given by Geramita and Gimigliano ([5]).

**Lemma 2.2.**  $\mathbf{E}$  has maximal rank.

*Proof.* We assume, without loss of generality, that none of the points of  $\mathbb{X}$  is  $P = [0 : 0 : 1]$ , and that the first minor of  $\mathbf{L}$ ,  $F_1$ , does not vanish at  $P$ . Suppose

$$\mathbf{L} = A_1 w_1 + A_2 w_2 + A_3 w_3,$$

where the  $A_i$ s have entries in the ground field. This means that  $A_3$  has maximal rank  $d$  (since  $F_1(P) \neq 0$ ).



$x_{1j}, x_{2j}, \dots, x_{uj}$  satisfy the  $2 \times 2$  minors of the Catalecticant matrix  $\text{Cat}(1, n-1; 3)$  of size  $3 \times \binom{n+1}{2}$ , for all  $j = 1, 2, \dots, d+1$ . Denote the collection of these equations by (\*\*\*) .

From (2.1), on  $\varphi(\mathbb{P}^2 \setminus \mathbb{X})$ , we have:

$$x_{1j}/z_1 = x_{2j}/z_2 = \dots = x_{uj}/z_u, \quad \text{for all } j = 1, 2, \dots, d+1.$$

This can be rewritten as a number of systems of equations, one for each  $i = 1, 2, \dots, u$

$$\left\{ \begin{array}{l} x_{ij}/z_i = x_{1j}/z_1 \\ x_{ij}/z_i = x_{2j}/z_2 \\ \dots \\ x_{ij}/z_i = x_{uj}/z_u \end{array} \right. \quad \text{for } j = 1, 2, \dots, d+1.$$

Those relations give us, for each  $i = 1, 2, \dots, u$  :

$$(S_i) \left\{ \begin{array}{l} x_{ij}z_1 - x_{1j}z_i = 0 \\ x_{ij}z_2 - x_{2j}z_i = 0 \\ \dots \\ x_{ij}z_u - x_{uj}z_i = 0 \end{array} \right. \quad \text{for } j = 1, 2, \dots, d+1.$$

It is not hard to see that if the coordinates of  $Q = [x_{ij}] \in \mathbb{P}^p$  and  $P = [z_i] \in \mathbb{P}^{u-1}$  satisfy system  $(S_i)$  for some  $i$ , where  $z_i \neq 0$ , then they satisfy systems  $(S_i)$  for all  $i$ .

Before going further, we prove a similar proposition to that of [5].

**Proposition 2.3.** *Let  $Q = [x_{ij}]$  be a point on  $\mathbb{P}^p$ , and suppose the coordinates of  $Q$  satisfy equations in (\*\*). Then there exists a unique  $P = [z_1 : \dots : z_u] \in \mathbb{P}^{u-1}$  such that the homogeneous coordinates of  $P$  and  $Q$  satisfy the systems  $(S_i)$  for all  $i$ .*

*Proof.* Since the coordinates of  $Q$  satisfy equations in (\*\*), the matrix  $M(Q)$  has rank 1, i.e. the rows of  $M(Q)$  are all multiples of any nonzero row of  $M(Q)$ . Suppose the first row of  $M(Q)$  is not identically zero (similar argument works for other rows). Then there exist  $\nu_i$ , for  $i = 2, \dots, u$ , such that

$$x_{ij} = \nu_i x_{1j}, \quad \text{for all } j = 1, 2, \dots, d+1.$$

We want  $P \in \mathbb{P}^{u-1}$  such that the coordinates of  $P$  and  $Q$  satisfy the systems  $(S_i)$  for all  $i$ . We first consider such  $P$  that the coordinates of  $P$  and  $Q$  satisfy  $(S_1)$ . This is the same as solving for  $z_1, \dots, z_u$  from  $(S_1)$ . The coefficients matrix becomes (projectively) a collection

of :

$$N_j = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 \\ -\nu_2 & 1 & 0 & \dots & 0 \\ -\nu_3 & 0 & 1 & \dots & 0 \\ \vdots & & & \ddots & \\ -\nu_u & 0 & 0 & \dots & 1 \end{pmatrix}, \text{ for } j = 1, 2, \dots, d+1.$$

Since  $N_j$  is independent of  $j$  and has rank exactly  $u - 1$ , the system  $(S_1)$  has exactly one projective solution, that gives a unique point  $P \in \mathbb{P}^{u-1}$ . Moreover, this  $P$  clearly has non-zero  $z_1$  entry. Thus, the coordinates of  $P$  and  $Q$  satisfy  $(S_i)$  for all  $i$ . Hence,  $P$  exists and is unique.  $\square$

Let  $\mathbf{V}$  be the algebraic set in  $\mathbb{P}^p$  defined by all the equations in (2.2), (\*\*) and (\*\*\*). We have the following theorem.

**Theorem 2.4.**  $\mathbf{V} = \overline{\Lambda_t}$  as sets.

*Proof.* Clearly,  $\varphi(\mathbb{P}^2 \setminus \mathbb{X}) \subseteq \mathbf{V}$ . Since  $\mathbf{V}$  is closed,  $\overline{\Lambda_t}$  is integral (so  $\overline{\Lambda_t}$  is irreducible, and  $\overline{\Lambda_t} = \overline{\varphi(\mathbb{P}^2 \setminus \mathbb{X})}$ ), we have

$$\overline{\Lambda_t} \subseteq \mathbf{V}.$$

We only need to show that

$$\mathbf{V} \subseteq \overline{\Lambda_t}.$$

Having Proposition 2.3, if we can show that for any points  $P = [z_1 : \dots : z_u]$  and  $Q = [x_{ij}]$  such that the coordinates of  $Q$  satisfy the equations in (2.2), (\*\*) and (\*\*\*), and coordinates of  $P$  and  $Q$  satisfy the systems  $(S_i)$  for all  $i$ ,  $Q$  must be in  $\overline{\Lambda_t}$ , then we will have  $\mathbf{V} \subseteq \overline{\Lambda_t}$ , and so are done. Suppose  $P$  and  $Q$  are such points. We can always assume that  $z_1 \neq 0$ . Consider the system of equations given by all the equations in  $(S_1)$  (if instead,  $z_i \neq 0$ , then we look at the system  $(S_i)$ ). As a system of linear equations in indeterminates (note the way we have rearranged the indices)

$$\{x_{ij} | 1 \leq j \leq d+1, 1 \leq i \leq u\},$$

the coefficients matrix is :

$$A = \begin{pmatrix} B & & & & \\ & B & & & \\ & & \ddots & & \\ & & & & B \end{pmatrix},$$

where

$$B = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 \\ z_2 & -z_1 & 0 & \dots & 0 \\ z_3 & 0 & -z_1 & \dots & 0 \\ \vdots & & & \ddots & \\ z_u & 0 & 0 & \dots & -z_1 \end{pmatrix}.$$

Clearly,  $B$  has rank  $u - 1$ , and has a non-trivial solution  $[z_1 : \dots : z_u]$ . Therefore, the solution to  $A$  must have the form :

$$[x_{ij}] = [c_1 z_1 : c_1 z_2 : \dots : c_1 z_u : c_2 z_1 : \dots : c_2 z_u : \dots : c_{d+1} z_1 : \dots : c_{d+1} z_u],$$

where  $c_1, \dots, c_{d+1}$  are constants not all zero, and the indeterminates are ordered by  $1 \leq j \leq d + 1$  and  $1 \leq i \leq u$ .

Now, since the coordinates of  $Q$  also satisfy the equations in (\*\*\*), which are the defining equations of Veronese surfaces, there exists a unique point  $T = [w_1 : w_2 : w_3] \in \mathbb{P}^2$  such that  $z_1 = w_1^n, z_2 = w_1^{n-1} w_2, \dots, z_u = w_3^n$ . Thus,

$$Q = [c_1 w_1^n : \dots : c_{d+1} w_3^n]. \quad (2.3)$$

Lastly, the coordinates of  $Q$  satisfy  $\binom{n+1}{2}d$  equations in (2.2), so

$$\mathbf{L}(T) \begin{bmatrix} c_1 \\ c_2 \\ \dots \\ c_{d+1} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \dots \\ 0 \end{bmatrix}.$$

If  $T \notin \mathbb{X}$ , then  $\mathbf{L}(T)$  has rank exactly  $d$ . Thus,

$$\begin{bmatrix} c_1 \\ c_2 \\ \dots \\ c_{d+1} \end{bmatrix} = \rho \begin{bmatrix} F_1(T) \\ F_2(T) \\ \dots \\ F_{d+1}(T) \end{bmatrix}. \quad (2.4)$$

This implies that  $Q \in \overline{\Lambda}_t$ .

If  $T \in \mathbb{X}$ , then  $\mathbf{L}(T)$  has rank exactly  $d - 1$ , so there is a 2-dimensional solution space, and these resulting  $Q$ s lie on a line of  $\mathbf{V}$ , which is one of the exceptional lines of  $\overline{\Lambda}_t$ .

Hence, we always have  $Q \in \overline{\Lambda}_t$ . We have proved that  $\mathbf{V} = \overline{\Lambda}_t$  as sets. □

To continue our study, we let  $S = \mathfrak{k}[x_{ij}]$  be the homogeneous coordinate ring of  $\mathbb{P}^p$ . Suppose  $\mathbf{C}$  is the Catalecticant matrix  $\text{Cat}(1, n - 1; 3)$  of indeterminates  $\{z_i\}_{0 \leq i \leq \binom{n+2}{2}}$ .  $\mathbf{C}$  is of size  $3 \times \binom{n+1}{2}$ . Consider the box  $\mathbf{B}$  of size  $(d + 1) \times 3 \times \binom{n+1}{2}$ . Let  $\mathcal{A}$  be the box-shaped matrix obtained by assigning each integral point  $(i, j, k)$  of  $\mathbf{B}$  the indeterminate  $x_{il}$  where  $l$  is the integer such that  $z_l$  is at the  $(j, k)$ -position in  $\mathbf{C}$ .

**Lemma 2.5.**  *$\mathcal{A}$  is a weak box-shaped matrix of indeterminates.*

*Proof.* Clearly, each x-section of  $\mathcal{A}$  has its ideal of  $2 \times 2$  minors as the defining ideal of a Veronese surface, so its ideal of  $2 \times 2$  minors is a prime ideal. Also, each y-section and z-section of  $\mathcal{A}$  is a matrix of indeterminates, whence whose ideal of  $2 \times 2$  minors is also a prime ideal. Moreover,  $x_{111}$  surely satisfies property (c) of  $\mathcal{A}$  being a weak box-shaped matrix. It remains to show that

$$\langle I_2(\mathcal{A}), x_{(d+1)3\binom{n+1}{2}} \rangle = \bigcap_{l=1}^3 I_l.$$

For convenience, we let  $r_1 = d + 1, r_2 = 3, r_3 = \binom{n+1}{2}$ , and consider  $\mathcal{A}$  as a box-shaped matrix of size  $r_1 \times r_2 \times r_3$ . We shall first prove that

$$I_2 \cap I_3 = \langle I_2(\mathcal{A}), \{x_{ir_2r_3} \mid i = 1, \dots, r_1\} \rangle.$$

The proof will go in the same line as that of part (a) of Lemma 1.1. Firstly, it is clear that  $\langle I_2(\mathcal{A}), \{x_{ir_2r_3} \mid i = 1, \dots, r_1\} \rangle \subseteq I_2 \cap I_3$ . It remains to show the other inclusion. Let  $F \in I_2 \cap I_3$ . Doing exactly as we did before, we end up with  $F = F' + G' + G$ , where  $F', G' \in I_2(\mathcal{A})$ , and

$$G = \sum_{ik} G_{ik} x_{ir_2k},$$

where  $G_{ik}$ 's are independent of the variables  $x_{ijr_3}$ . Again, we have  $G \in I_3$ , so we can write

$$G = H + \sum_{i,j} H_{ij} x_{ijr_3},$$

where  $H \in I_2(\mathcal{A}_3)$ . We may assume that the  $H_{ir_2}$ 's are independent of all the variables  $\{x_{ijr_3} | j \neq r_2\}$ . By the nature of the  $2 \times 2$  minors of  $\mathcal{A}$  and the symmetry (in construction) of Catalecticant matrices, it can be seen that if a  $2 \times 2$  minor of  $\mathcal{A}$  has one indeterminate belonging to  $\{x_{ijr_3} | (i, j, r_3) \in \mathbf{B}\}$  then it must have at least two adjacent indeterminates belonging to  $\{x_{ijr_3} | (i, j, r_3) \in \mathbf{B}\}$ . Thus, by re-grouping and rewriting, we can always assume that  $H$  is also independent of the indeterminates  $\{x_{ijr_3} | (i, j, r_3) \in \mathbf{B}\}$ . Now, clearly,  $G = H + \sum_i H_{ir_2} x_{ir_2r_3} \in \langle I_2(\mathcal{A}), \{x_{ir_2r_3} | i = 1, \dots, r_1\} \rangle$ . We have shown that  $I_2 \cap I_3 = \langle I_2(\mathcal{A}), \{x_{ir_2r_3} | i = 1, \dots, r_1\} \rangle$ .

It now follows in the same line of the proof of part (b) of Lemma 1.1 that

$$\langle I_2(\mathcal{A}), \{x_{ir_2r_3} | i = 1, \dots, r_1\} \rangle \cap I_3 = \langle I_2(\mathcal{A}), x_{r_1r_2r_3} \rangle .$$

The lemma is proved. □

We obtain the main result of this section as follows.

**Theorem 2.6.** *The subscheme  $\overline{\Lambda}_t$  in  $\mathbb{P}^p$  is defined by  $\binom{n+1}{2}d$  linear forms and the  $2 \times 2$  minors of a box-shaped matrix of linear forms.*

*Proof.* Let  $S = \mathfrak{k}[x_{ij}]$  be the homogeneous coordinate ring of  $\mathbb{P}^p$ . Let  $\mathcal{A}$  be the weak box-shaped matrix of indeterminates as above, and again, let  $I_2(\mathcal{A})$  be the ideal of  $2 \times 2$  minors of  $\mathcal{A}$  in  $\mathfrak{k}[\mathcal{A}]$ . We also let  $\mathbf{I}$  be the ideal generated by  $I_2(\mathcal{A})$  and all the linear equations in (2.2). Let  $\mathcal{V}$  be the subscheme of  $\mathbb{P}^p$  defined by  $\mathbf{I}$ .

It is easy to see that  $\mathbf{I}$  contains all the equations in (2.2), (\*\*), and (\*\*\*), so as sets,  $\mathcal{V} \subseteq \mathbf{V}$  (where  $\mathbf{V}$  is the subvariety of  $\mathbb{P}^p$  defined by the equations in (2.2), (\*\*), and (\*\*\*)).

Suppose now that  $P = [\overline{w}_1 : \overline{w}_2 : \overline{w}_3] \in \mathbb{P}^2 \setminus \mathbb{X}$  and  $Q = [\overline{x}_{ij}] = \varphi(P)$ . Let  $\overline{z}_1 = \overline{w}_1^n, \overline{z}_2 = \overline{w}_1^{n-1}\overline{w}_2, \dots, \overline{z}_u = \overline{w}_3^n$  then  $\overline{x}_{ij} = \overline{z}_i F_j(P)$ . Consider a  $2 \times 2$  minors  $a_{(K,L,M,N)}$  of  $\mathcal{A}$  corresponding to the 4 points  $K, L, M$  and  $N$  in the box-shaped realization of  $\mathcal{A}$ . There are 3 possibilities for the tuple  $(K, L, M, N)$ .

**Case 1.**  $K = (i, j, k), L = (m, j, p), M = (m, n, p)$  and  $N = (i, n, k)$  for some integers  $i, j, k, m, n$  and  $p$  (when the projections of  $K, L, M, N$  on the  $zx$ -plane collapse to a line).

**Case 2.**  $K = (i, j, k), L = (m, j, k), M = (m, n, p)$  and  $N = (i, n, p)$  for some integers  $i, j, k, m, n$  and  $p$  (when the projections of  $K, L, M, N$  on the  $yz$ -plane collapse to a line).

**Case 3.**  $K = (i, j, k), L = (m, n, k), M = (m, n, p)$  and  $N = (i, j, p)$  for some integers  $i, j, k, m, n$ , and  $p$  (when the projections of  $K, L, M, N$  on the  $xy$ -plane collapse to a line).

By the construction of  $\mathcal{A}$  and the fact that  $[\bar{z}_1 : \dots : \bar{z}_u]$  is in the Veronese surface, i.e. it satisfies all the  $2 \times 2$  minors of  $\mathbf{C}$ , it is easy to check that  $Q$  satisfies the minors  $a_{(K,L,M,N)}$ . This is true for any  $Q \in \varphi(\mathbb{P}^2 \setminus \mathbb{X})$  and any  $2 \times 2$  minors  $a_{(K,L,M,N)}$  of  $\mathcal{A}$ , so  $\varphi(\mathbb{P}^2 \setminus \mathbb{X}) \subseteq \mathcal{V}$ , whence  $\mathbf{V} \subseteq \mathcal{V}$ .

We have shown that in all cases,  $\mathbf{V} \subseteq \mathcal{V}$ . Hence, as sets,  $\mathcal{V} = \mathbf{V} = \overline{\Lambda}_t$ .

Now, by Proposition 1.12, we know that  $I_2(\mathcal{A})$  is a prime ideal. Consider the following sequence of surjective ring homomorphisms:

$$\mathfrak{k}[x_{ij}] \xrightarrow{\phi} \mathfrak{k}[w^\alpha t_j] \xrightarrow{\psi} \mathfrak{k}[w^\alpha F_j],$$

defined in the obvious way; that is, both  $\phi$  and  $\psi$  send  $\mathfrak{k}$  to  $\mathfrak{k}$ , and  $\phi$  sends  $x_{ij}$  to  $w^\alpha t_j$  where  $w^\alpha$  is labelled  $z_i$ , and  $\psi$  sends  $w^\alpha t_j$  to  $w^\alpha F_j$ .

We note that in proving equalities (2.3) and (2.4), we actually proved more. Firstly, the proof of (2.3) and the fact that  $I_2(\mathcal{A})$  is a prime ideal imply that  $I_2(\mathcal{A})$  is the kernel of  $\phi$ . Secondly, the proof of (2.4) shows that if we consider the equations in (2.2) as polynomials over the  $w^\alpha t_j$ 's, then those polynomials are zero exactly when  $t_j = F_j$  (since  $t_j = F_j$  at all but a finite set of points  $\mathbb{X}$ ). This implies that  $\mathfrak{k}[w^\alpha t_j]/\mathfrak{a} \simeq \mathfrak{k}[w^\alpha F_j]$ , where  $\mathfrak{a}$  is the ideal generated by the images of the equations in (2.2) through  $\phi$ . Thus,  $\mathfrak{a}$  is the kernel of  $\psi$ . Now, by Lemma 2.1, we conclude that  $\mathbf{I}$  is the kernel of  $\psi \circ \phi$ . In other words,  $\mathbf{I}$  is the defining ideal of  $\overline{\Lambda}_t$  embedded in  $\mathbb{P}^p$  (since the homogeneous coordinate ring of  $\overline{\Lambda}_t$  embedded in  $\mathbb{P}^p$  is exactly  $\mathfrak{k}[w^\alpha F_j]$ ). The theorem is proved.  $\square$

**Remark:** When  $t = d + 1$ , our box-shaped matrix  $\mathcal{A}$  collapses to be a normal matrix of size  $3 \times (d + 1)$ , and the above result coincides with that obtained by Geramita and Gimigliano in [5].

*Acknowledgement.* This paper is part of the author's PhD thesis. I would like to thank my research advisor A.V. Geramita for his inspiration and guidance.

## REFERENCES

- [1] Ciliberto, C., Geramita, A.V. and Orecchia, F. (1988). *Remarks on a theorem of Hilbert-Burch*. Boll. Unione. Math. Ital. **7**, 2-B, 463-483.
- [2] Conca, A., Herzog, J., Trung, N.V., and Valla, G. (1997). *Diagonal subalgebras of bigraded algebras and embeddings of blow-ups of projective spaces*. Amer. J. Maths., **119**, 859-901.
- [3] Davis, E.D. and Geramita, A.V. (1988). *Birational morphisms to  $\mathbb{P}^2$ : an ideal-theoretic perspective*. Math. Ann. **83**, 435-448.
- [4] Davis, E.D., Geramita, A.V. and Maroscia, P. (1984). *Perfect homogeneous ideals: Dubreil's theorems revisited*. Bull. Sci. Math. **108**, 143-185.
- [5] Geramita, A.V. and Gimigliano, A. (1991). *Generators for the defining ideal of certain rational surfaces*. Duke Mathematical Journal, **62**, no. 1, 61-83.
- [6] Geramita, A.V., Gimigliano, A. and Harbourne, B. (1994). *Projectively normal but superabundant embeddings of rational surfaces in projective space*. J. Algebra. **169**, no 3, 791-804.
- [7] Geramita, A.V., Gimigliano, A. and Pitteloud, Y. (1995). *Graded Betti numbers of some embedded rational  $n$ -folds*. Math. Ann. **301**, 363-380.
- [8] Geramita, A.V., Gregory, D. and Roberts, L.G. (1986). *Monomial ideals and points in projective space*. J. Pure Appl. Algebra. **40**, 33-62.
- [9] Geramita, A.V. and Maroscia, P. (1984). *The ideal of forms vanishing at a finite set of points in  $\mathbb{P}^n$* . Journal of Algebra, **90**, 528-555.
- [10] Gimigliano, A. (1987). *Linear systems of plane curves*. PhD Thesis. Queen's University, Kingston.
- [11] Gimigliano, A. (1989). *On Veronesean surfaces*. Proc. Konin. Ned. Akad. van Wetenschappen, Ser. A, **92**, 71-85.
- [12] Gimigliano, A. and Loenzini, A. (1993). *On the ideal of veronesean surfaces*. Can. J. Math. **43**, 758-777.
- [13] Grone, Robert (1977). *Decomposable tensors as a quadratic variety*. Proc. of Amer. Math. Soc. **64**, no 2, 227-230.
- [14] Hà, Huy Tàì (2000). *Rational surfaces from an algebraic perspective*. PhD Thesis. Queen's University, Kingston.
- [15] Harbourne, B. (1986). *The geometry of rational surfaces and Hilbert functions of points in the plane*. Canad. Math. Soc. Conf. Proc. **6**, 95-111.
- [16] Hochster, M. and Eagon, A. (1971). *Cohen-Macaulay rings, invariant theory, and the generic perfection of determinantal loci*. Amer. J. Math. **93**, 1020-1058.

- [17] Hochster, M. (1972). *Rings of invariant of tori, Cohen-Macaulay rings generated by monomials, and polytopes*. Ann. of Math. (2). **96**, 318-337.
- [18] Holay, S.H. (1994). *Generators and resolutions of ideals defining certain surfaces in projective space*. PhD Thesis. University of Nebraska, Lincoln.
- [19] Holay, S.H. (1996). *Generators of ideals defining certain surfaces in projective space*. Canad. J. Math. **48**, no 3, 585-595.
- [20] Holay, S.H. (1996). *Free resolutions of the defining ideal of certain rational surfaces*. Manuscripta Math. **90**, no. 1, 23-37.
- [21] Huneke, C. (1981). *The arithmetic perfection of Buchsbaum-Eisenbud varieties and generic modules of projective dimension two*. Trans. Amer. Math. Soc. **265**. 211-233.
- [22] Morey, S. and Ulrich, B. (1996). *Rees algebras of ideals with low codimension*. Proc. Amer. Math. Soc. **124**, no. 12, 3653-3661.
- [23] Morey, S. (1996). *Equations of blowups of ideals of codimension two and three*. J. Pure Appl. Algebra, **109**, 197-211.
- [24] Room, T.G. (1938). *The Geometry of Determinantal Loci*. Cambridge University Press, Cambridge.
- [25] Pucci, M. (1998). *The Veronese variety and Catalecticant matrices*. J. Algebra. **202**, no. 1, 72-95.
- [26] Sharpe, D.W. (1964). *On certain polynomial ideals defined by matrices*. Quart. J. Math. Oxford (2), **15**, 155-175.
- [27] Simis, A., Trung, N.V. and Valla, G. (1998). *The diagonal subalgebra of a blow-up algebra*. J. Pure Appl. Algebra, **125**, 305-328.
- [28] Stückrad, J. and Vogel, W. (1978). *On Segre Products and Applications*. J. Algebra. **54**, 374-389.
- [29] Trung, N.V. and Valla, G. (1989). *The Cohen-Macaulay type of points in generic position*. J. Algebra. **125**, 110-119.
- [30] White, M.P. (1922). *The projective generation of curves and surfaces in space of four dimensions*. Proc. Cambridge Phil. Soc., **21**, 216-227.

DEPARTMENT OF MATHEMATICS AND STATISTICS, QUEEN'S UNIVERSITY, KINGSTON ON K7L 3N6, CANADA  
*E-mail address:* haht@mast.queensu.ca