

ASYMPTOTIC BEHAVIOR OF THE LENGTH OF LOCAL COHOMOLOGY

STEVEN DALE CUTKOSKY, HUY TÀI HÀ, HEMA SRINIVASAN, AND EMANOIL THEODORESCU

ABSTRACT. Let k be a field of characteristic 0, $R = k[x_1, \dots, x_d]$ be a polynomial ring, and \mathfrak{m} its maximal homogeneous ideal. Let $I \subset R$ be a homogeneous ideal in R . In this paper, we show that

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d} = \lim_{n \rightarrow \infty} \frac{\lambda(\text{Ext}_R^d(R/I^n, R(-d)))}{n^d}$$

always exists. This limit has been shown to be $\frac{e(I)}{d!}$ for \mathfrak{m} -primary ideals I in a local Cohen Macaulay ring [Ki, Ko1, Th, Th2], where $e(I)$ denotes the multiplicity of I . But we find that this limit may not be rational in general. We give an example for which the limit is an irrational number thereby showing that the lengths of these extension modules may not have polynomial growth.

INTRODUCTION

Let $R = k[x_1, \dots, x_d]$ be a polynomial ring over a field k , with graded maximal ideal \mathfrak{m} , and $I \subset R$ a proper homogeneous ideal. We investigate the asymptotic growth of $\lambda(\text{Ext}_R^d(R/I^n, R))$ as a function of n . When R is a local Gorenstein ring and I is an \mathfrak{m} -primary ideal, then this is easily seen to be equal to $\lambda(R/I^n)$ and hence is a polynomial in n . A theorem of Kirby, Kodiyalam and Theodorescu [Ki, Ko1, Th, Th2] extends this to \mathfrak{m} -primary ideals in local Cohen Macaulay rings R . We consider homogeneous ideals in a polynomial ring which are not \mathfrak{m} -primary and show that a limit exists asymptotically although it can be irrational. In our setting, by local duality,

$$\lambda(\text{Ext}_R^d(R/I^n, R(-d))) = \lambda(H_{\mathfrak{m}}^0(R/I^n))$$

and thus this becomes a problem of asymptotic lengths of local cohomology modules.

In recent years, a great deal of interest has been given to investigating asymptotic behavior of algebraic invariants of powers of I . Cutkosky, Herzog and Trung [CHT], and Kodiyalam [Ko2] independently proved that $\text{reg}(R/I^n)$ is a linear function in n for $n \gg 0$ (see also [Ch, GGP]). When I^n is replaced by its saturation $(I^n)^{\text{sat}}$, the problem becomes much subtler. It is no longer true that $\text{reg}(R/(I^n)^{\text{sat}})$ is always asymptotically a polynomial in n as shown in [Cu]. Examples are given in [Cu, CEL] showing that it is possible for $\lim_{n \rightarrow \infty} \frac{\text{reg}(R/(I^n)^{\text{sat}})}{n}$ to be an irrational number. Further, Cutkosky, Ein and Lazarsfeld [CEL] showed that the limit $\lim_{n \rightarrow \infty} \frac{\text{reg}(R/(I^n)^{\text{sat}})}{n}$ always exists. Along this theme, Hoa and Hyry [HoH] recently studied the existence of similar limits where the regularity of R/I^n is replaced by its a-invariants. This paper addresses a closely related question. We prove

1991 *Mathematics Subject Classification.* 13D40, 14B15, 13D45.

Key words and phrases. powers of ideals, local cohomology, Hilbert function, linear growth.

The first author was partially supported by NSF.

The third author was partially supported by NSA.

Theorem 0.1. *Let k be a field of characteristic zero, $R = k[x_1, \dots, x_d]$ be a polynomial ring of dimension $d > 1$, and \mathfrak{m} its maximal homogeneous ideal. Let $I \subset R$ be a homogeneous ideal of R . Then, the limit*

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d} = \lim_{n \rightarrow \infty} \frac{\lambda(\text{Ext}_R^d(R/I^n, R(-d)))}{n^d}$$

always exists.

In fact, we prove (in Theorem 1.3) that if R is a coordinate ring of a projective variety which has depth ≥ 2 at its irrelevant ideal, then the limit $\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d}$ exists, and (Corollary 1.4) if R is Gorenstein, then

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d} = \lim_{n \rightarrow \infty} \frac{\lambda(\text{Ext}_R^d(R/I^n, R(-d)))}{n^d}.$$

We will also give an example where $\frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d}$ tends to an irrational number as $n \rightarrow \infty$ (Theorem 2.2). This, in particular, shows that, just like $\text{reg}(R/(I^n)^{\text{sat}})$, $\lambda(H_{\mathfrak{m}}^0(R/I^n))$ is not asymptotically a polynomial in n .

Theorem 0.1 is proved in Section 1. To do this, we express $\lambda(H_{\mathfrak{m}}^0(R/I^n))$ as a sum of two components, the geometric component $\sigma(n)$, and the algebraic component $\tau(n)$, and show that both limits $\lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^d}$ and $\lim_{n \rightarrow \infty} \frac{\tau(n)}{n^d}$ exist. For the first limit, we express $\sigma(n)$ as $h^0(Y, \mathcal{N}^n)$ for some line bundle \mathcal{N} over a projective scheme Y of dimension d , and investigate the limit $\lim_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{N}^n)}{n^d}$. For the later one, we write $\tau(n)$ as the Hilbert function of a finitely generated graded k -algebra of dimension $(d+1)$. In Theorem 0.2, we use the construction illustrated in [Cu] by the first author to give an example where the limit proved to exist in Section 1 is an irrational number.

Theorem 0.2. *There exists a nonsingular projective curve $C \subset \mathbb{P}_{\mathbb{C}}^3$ such that if $I \subset R = \mathbb{C}[x_1, \dots, x_4]$ is the defining ideal of C , and \mathfrak{m} is the homogeneous maximal ideal of R , then*

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^4} \notin \mathbb{Q}.$$

Of course, Theorem 0.2 has a local analog.

Theorem 0.3. *There exists a regular local ring S of dimension 4 which is essentially of finite type over the complex numbers \mathbb{C} , and an ideal $J \subset S$ such that*

$$\lim_{n \rightarrow \infty} \frac{\lambda(\text{Ext}_S^d(S/J^n, S))}{n^d}$$

is an irrational number. In particular, $\lambda(\text{Ext}_S^d(S/J^n, S))$ is not a polynomial or a quasi-polynomial for large n .

The proofs of Theorems 0.2 and 0.3 will be given in section 2. In contrast to the example of Theorem 0.3, if (S, \mathfrak{m}, k) is a Cohen-Macaulay local ring of dimension d and J is an \mathfrak{m} -primary ideal, then $\lambda(\text{Ext}_S^d(S/J^n, S))$ is a polynomial of degree d for large n ([Ki], [Th]). In fact,

$$\lim_{n \rightarrow \infty} \frac{\lambda(\text{Ext}_S^d(S/J^n, S))}{n^d} = \frac{e(I)}{d!}$$

where $e(I)$ is the multiplicity of I ([Th2]).

In the case when S is Gorenstein, this follows easily from local duality, since

$$\lambda(\text{Ext}^d(S/J^n, S)) = \lambda(H_{\mathfrak{m}_S}^0(S/J^n)) = \lambda(S/J^n)$$

for all n .

Suppose that I is a homogeneous ideal in the coordinate ring R of a projective variety of depth ≥ 2 at the irrelevant ideal. Then (c.f. Remark 1.5), if $\text{ht } I = d = \dim(R)$, we have that

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d} = \frac{e(I)}{d!}$$

where $e(I)$ is the multiplicity of I .

In contrast, if $\text{ht } I < d$ then

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d} \tag{0.1}$$

does not have such a simple arithmetic interpretation. The example of Theorem 0.2 of this paper is of a height 2 prime ideal in a polynomial ring R of dimension 4 such that (0.1) is an irrational number. However, in many cases, such as when I is a regular prime with $\text{ht } I < d$ in a polynomial ring R of dimension d , we have that the limit (0.1) is 0.

More generally, if $\text{ht } I < d$ and the analytic spread $\ell(I) < d$ then the integral closure $\overline{I^n}$ has no \mathfrak{m} -primary component for large n [M, Theorem 3]. Since $\text{depth}(R_{\mathfrak{m}}) \geq 2$, $\overline{I^n} = H^0(\text{spec}(R) - \{\mathfrak{m}\}, \overline{I^n})$, and

$$H_{\mathfrak{m}}^0(R/\overline{I^n}) \cong H_{\mathfrak{m}}^1(\overline{I^n}) \cong H^0(\text{spec}(R) - \{\mathfrak{m}\}, \overline{I^n}/\overline{I^n}) = 0$$

for large n . Thus if I is a normal ideal ($I^n = \overline{I^n}$ for all n) with analytic spread $\ell(I) < d$ we have that $H_{\mathfrak{m}}^0(R/I^n) = 0$ for large n and the limit (0.1) is thus 0. In fact, Catalin Ciuperca has shown us that even if I is not normal, with $\ell(I) < d$, then the limit (0.1) is zero.

1. THE EXISTENCE THEOREM

In this section, we prove the main theorem of the paper. We shall start by recalling some notations and terminology, and prove a few preliminary results.

Suppose R is a graded ring, and $I \subset R$ a homogeneous ideal. The *Rees algebra* of I is the subalgebra $R[It]$ of $R[t]$. The Rees algebra $R[It]$ has a natural bi-gradation given by

$$R[It]_{(m,n)} = (I^n)_m t^n.$$

Suppose $A = \bigoplus_{m,n \in \mathbb{Z}} A_{(m,n)}$ is a bi-graded algebra. For a tuple of positive integers $\Delta = (a, b)$, $A_{\Delta} = \bigoplus_{n \in \mathbb{Z}} A_{(an, bn)}$ is call a Δ -*diagonal subalgebra* of A .

Lemma 1.1. *Suppose a domain R is a finitely generated graded k -algebra of dimension δ , and $I \subset R$ is a homogeneous ideal generated in degrees $\leq d$ such that $\text{ht } I \geq 1$. Let $A = R[It]$ be the Rees algebra of I over R .*

(i) *For any tuple $\Delta = (a, b)$ of positive integers such that $a \geq db$, A_{Δ} is a finitely generated graded k -algebra.*

(ii) For any tuple $\Delta = (a, b)$ of positive integers such that $a > db$, $\dim A_\Delta = \delta$.

Proof. It is easy to see that $A_\Delta = k[(I^b)_a]$ is the k -algebra generated by elements of $(I^b)_a$. Thus, (i) is clear. (ii) follows from [HgT, Lemma 2.2] since R is a domain. \square

The following Lemma is stated in an example in [La].

Lemma 1.2. *Suppose that Y is a projective variety of dimension d over a field k of characteristic zero, and \mathcal{L} is a line bundle on Y . Then, the limit*

$$\lim_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d}$$

exists, and is a positive real number if \mathcal{L} is big.

Proof. It follows from [I, Theorem 10.2] that $\limsup_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d} = 0$ if \mathcal{L} is not big ($\kappa(\mathcal{L}) < d$). This implies that $\lim_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d} = 0$.

Suppose that \mathcal{L} is big (that is, $\kappa(\mathcal{L}) = d$). It follows from [I, Theorem 10.2] that

$$\liminf_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d} > 0.$$

To prove the lemma, it suffices to show that

$$\limsup_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d/d!} = \liminf_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d/d!}.$$

Let $\varepsilon > 0$ be an arbitrary positive number. By applying the theorem of Fujita [Fu] (and from the definition of \limsup), there exists a birational morphism $\theta : Z \rightarrow Y$ together with an effective \mathbb{Q} -divisor E on Z such that $H = \theta^*\mathcal{L} - E$ is a semiample \mathbb{Q} -divisor with

$$H^d > \limsup_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d/d!} - \varepsilon.$$

Let q be the smallest positive integer such that qE is integral (or equivalently, qH is integral). Since qE is effective, there is a natural injection $\mathcal{O}_Z \hookrightarrow \mathcal{O}_Z(qE)$. This gives an injection $\mathcal{O}_Z(lqH) \hookrightarrow \mathcal{O}_Z(lqE + lqH) = \theta^*\mathcal{L}^{lq}$ for any integer $l > 0$. Thus, $h^0(Z, \mathcal{O}_Z(lqH)) \leq h^0(Z, \theta^*\mathcal{L}^{lq})$ for any integer $l > 0$. Furthermore, since qH is semiample, there exists $c > 0$ such that

$$h^i(Z, \mathcal{O}_Z(lqH)) \leq c(l^{d-2}) \tag{1.1}$$

for any $i \geq 1$ (c.f. Corollary 6.7 [F2]). Thus

$$h^0(Z, \mathcal{O}_Z(lqH)) = \chi(lqH) + O(l^{d-2}) = \frac{(lq)^d}{d!} H^d + O(l^{d-1}),$$

where χ denotes the Euler characteristic.

We have an exact sequence of coherent \mathcal{O}_Y modules

$$0 \rightarrow \mathcal{O}_Y \rightarrow \theta_*\mathcal{O}_Z \rightarrow \mathcal{F} \rightarrow 0$$

where \mathcal{F} is supported on a closed subset of Y of dimension $< d$. From the exact sequences

$$0 \rightarrow \mathcal{L}^{lq} \rightarrow \theta_*\theta^*\mathcal{L}^{lq} \rightarrow \mathcal{F} \otimes \mathcal{L}^{lq} \rightarrow 0$$

we have

$$h^0(Y, \mathcal{L}^{lq}) = h^0(Z, \theta^* \mathcal{L}^{lq}) + O(l^{d-1}).$$

Hence,

$$H^d = \lim_{l \rightarrow \infty} \frac{h^0(Z, \mathcal{O}_Z(lqH))}{(lq)^d/d!} \leq \liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq})}{(lq)^d/d!}.$$

In summary, we have

$$\limsup_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d/d!} < H^d + \varepsilon \leq \liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq})}{(lq)^d/d!} + \varepsilon. \quad (1.2)$$

Since ε is taken to be arbitrary, by (1.2), to prove the lemma, we only need to show that

$$\liminf_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d/d!} = \liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq})}{(lq)^d/d!}. \quad (1.3)$$

Since \mathcal{L} is big, there exists a constant n_0 such that $h^0(Y, \mathcal{L}^n) > 0$ for any $n \geq n_0$ (as follows from [I, Theorem 10.2]). This implies that, for each $0 \leq r < q$, there exists an effective divisor F_r on Y such that $\mathcal{O}_Y(F_r) \cong \mathcal{L}^{n_0+r}$. For $l > 0$, we have exact sequences

$$0 \longrightarrow \mathcal{L}^{lq} \longrightarrow \mathcal{L}^{lq+n_0+r} \longrightarrow \mathcal{O}_{F_r} \otimes \mathcal{L}^{lq+n_0+r} \longrightarrow 0.$$

Taking the long exact sequences of cohomologies, we get

$$0 \rightarrow H^0(Y, \mathcal{L}^{lq}) \rightarrow H^0(Y, \mathcal{L}^{lq+n_0+r}) \rightarrow H^0(F_r, \mathcal{O}_{F_r} \otimes \mathcal{L}^{lq+n_0+r}).$$

Thus,

$$\frac{h^0(Y, \mathcal{L}^{lq+n_0+r})}{(lq)^d/d!} - \frac{h^0(Y, \mathcal{L}^{lq+n_0+r})}{(lq)^d/d!} \leq \frac{h^0(F_r, \mathcal{O}_{F_r} \otimes \mathcal{L}^{lq+n_0+r})}{(lq)^d/d!}. \quad (1.4)$$

Since every component of F_r has dimension $d-1 < d$, we have

$$\lim_{l \rightarrow \infty} \frac{h^0(F_r, \mathcal{O}_{F_r} \otimes \mathcal{L}^{lq+n_0+r})}{(lq)^d/d!} = 0.$$

(1.4) now gives us

$$\liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq})}{(lq)^d/d!} = \liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq+n_0+r})}{(lq)^d/d!}.$$

Moreover, since n_0 is fixed and $0 \leq r < q$, we have

$$\liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq+n_0+r})}{(lq)^d/d!} = \liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq+n_0+r})}{(lq+n_0+r)^d/d!}.$$

Therefore,

$$\liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq})}{(lq)^d/d!} = \liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq+n_0+r})}{(lq+n_0+r)^d/d!}$$

for any $0 \leq r < q$. Hence,

$$\liminf_{l \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^{lq})}{(lq)^d/d!} = \liminf_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{L}^n)}{n^d/d!}.$$

(1.3) is proved, and so is the lemma. \square

Theorem 1.3. *Let $R = k[x_1, \dots, x_p]/J$ be the quotient of a polynomial ring $k[x_1, \dots, x_p]$ over a field k of characteristic zero by a homogeneous prime ideal J . Let \mathfrak{m} be the maximal homogeneous ideal of R . Suppose that $\text{depth}(R_{\mathfrak{m}}) \geq 2$. Let d be the dimension of R .*

Let $I \subset R$ be a homogeneous ideal of R . Then, the limit $\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d}$ always exists.

By duality, we obtain the following corollary.

Corollary 1.4. *With the notations of Theorem 1.3, suppose that R is Gorenstein. Then*

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^d} = \lim_{n \rightarrow \infty} \frac{\lambda(\text{Ext}_R^d(R/I^n, R(-d)))}{n^d}$$

Proof of Theorem 1.3: The problem is trivial if $\text{ht } I = 0$. If $\text{ht } I = d$ then there is a very simple proof (see the Remark after this proof). Suppose $\text{ht } I \geq 1$. Let \mathcal{I} be the ideal sheaf associated to I on $V = \text{Proj } R$. From the exact sequence

$$0 \longrightarrow I^n \longrightarrow R \longrightarrow R/I^n \longrightarrow 0,$$

we get $H_{\mathfrak{m}}^0(R/I^n) = H_{\mathfrak{m}}^1(I^n)$. The Serre-Grothendieck correspondence gives us the exact sequence

$$0 \rightarrow I^n \rightarrow \bigoplus_{m \geq 0} H^0(V, \mathcal{I}^n(m)) \rightarrow H_{\mathfrak{m}}^1(I^n) \rightarrow 0.$$

$\text{depth}(R_{\mathfrak{m}}) \geq 2$ implies that

$$\bigoplus_{m \geq 0} H^0(V, \mathcal{I}^n(m)) = H^0(\text{spec}(R) - \{\mathfrak{m}\}, I^n) = (I^n)^*$$

where $(I^n)^*$ is the intersection of the primary components of I^n which are not \mathfrak{m} -primary. By the theorem of Swanson [S] there exists a number $e > 0$ such that $(I^n)_m = (I^n)_m^*$ for any $m \geq en$ and $n \geq 1$. Therefore, we have

$$\lambda(H_{\mathfrak{m}}^1(I^n)) = \sigma(n) - \tau(n), \tag{1.5}$$

where

$$\sigma(n) = \sum_{m=0}^{en} h^0(V, \mathcal{I}^n(m)) \text{ and } \tau(n) = \sum_{m=0}^{en} \lambda((I^n)_m). \tag{1.6}$$

We will take e to be bigger than the degrees of homogeneous generators of I . The theorem will be proved if we can show that both limits $\lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^d}$ and $\lim_{n \rightarrow \infty} \frac{\tau(n)}{n^d}$ exist.

Let us first consider $\lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^d}$. Let $\pi : X \rightarrow V$ be the blowing up of V along \mathcal{I} . Let us denote $\mathcal{M} = \pi^* \mathcal{O}_V(1)$ and $\mathcal{L} = \mathcal{I} \mathcal{O}_X$. Let $\eta : Y = \mathbb{P}(\mathcal{O}_X \oplus \mathcal{M}) \rightarrow X$ be the projectivization of the vector bundle $\mathcal{O}_X \oplus \mathcal{M}$ on X . Then, $\dim Y = \dim X + 1 = d$. Let $\mathcal{N} = \mathcal{O}_Y(e) \otimes \eta^* \mathcal{L}$. We have

$$h^0(Y, \mathcal{N}^n) = h^0(X, S^{en}(\mathcal{O}_X \oplus \mathcal{M}) \otimes \mathcal{L}^n) = \sum_{m=0}^{en} h^0(X, \mathcal{M}^m \otimes \mathcal{L}^n).$$

Furthermore, it follows from [H, Exercise II.5.9] (see also [CEL, Lemma 3.3] and [HaT]) that $\pi_* \mathcal{L}^n = \mathcal{I}^n$ for $n \gg 0$. Thus, for $n \gg 0$,

$$h^0(Y, \mathcal{N}^n) = \sum_{m=0}^{en} h^0(V, \mathcal{O}_V(m) \otimes \mathcal{I}^n) = \sum_{m=0}^{en} h^0(V, \mathcal{I}^n(m)) = \sigma(n).$$

By Lemma 1.2, $\lim_{n \rightarrow \infty} \frac{h^0(Y, \mathcal{N}^n)}{n^d}$ exists. Hence, there exists a limit $\lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^d}$.

Now, let us consider $\lim_{n \rightarrow \infty} \frac{\tau(n)}{n^d}$. Suppose I is generated by $F_1, \dots, F_l \in R$ and $\deg F_j = d_j$ for $1 \leq j \leq l$. Let $S = R[s, F_1 t, \dots, F_l t] \subset R[s, t]$ be the Rees algebra of the ideal $IR[s]$ over the polynomial ring $R[s]$. S can be viewed as a bi-graded ring with $\deg x_i = (1, 0)$ for $1 \leq i \leq d$, $\deg s = (1, 0)$, and $\deg F_j t = (d_j, 1)$ for $1 \leq j \leq l$. Take an arbitrary element $f \in S$. We observe that $\deg f = (en, n)$ if and only if f has the following form

$$f = \sum_{m_1 + \dots + m_l = n} \sum_{j=0}^{en - d_1 m_1 - \dots - d_l m_l} b_j s^{en - d_1 m_1 - \dots - d_l m_l - j} F_1^{m_1} \dots F_l^{m_l} t^n,$$

where $b_j \in R$ is homogeneous of degree j . Thus the map

$$\Phi_n : S_{(en, n)} \rightarrow \sum_{m=0}^{en} (I^n)_m$$

defined by $\Phi_n(f(s, t)) = f(0, 1)$ is a k -vector space isomorphism. Hence,

$$\lambda(S_{(en, n)}) = \sum_{m=0}^{en} \lambda((I^n)_m) = \tau(n).$$

Let $T = \bigoplus_{n=0}^{\infty} S_{(en, n)}$, then $T = S_{\Delta}$ with $\Delta = (e, 1)$. Since S is a finitely generated bi-graded k -algebra and e is taken to be bigger than d_j for all $1 \leq j \leq l$, it follows from Lemma 1.1 that T is a finitely generated k -algebra and $\dim T = d + 1$. Thus, the Hilbert function $H(T, n) = \dim_k T_n = \lambda(S_{(en, n)})$ is given by a polynomial of degree d in n for $n \gg 0$ with a rational leading coefficient. This implies the existence of the limit

$$\lim_{n \rightarrow \infty} \frac{\tau(n)}{n^d} \in \mathbb{R}. \quad (1.7)$$

The theorem is proved.

Remark 1.5. Let assumptions be as in the statement of Theorem 1.3. If $\text{ht } I = d$ we have that $H_m^0(R/I^n) = R/I^n$ for all n , so that $\lambda(H_m^0(R/I^n))$ is the Hilbert polynomial of I for large n . Thus

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_m^0(R/I^n))}{n^d} = \frac{e(I)}{d!}$$

where $e(I)$ is the multiplicity of I . In the proof we in fact have that $\sigma(n) = \sum_{m=0}^{en} h^0(V, \mathcal{O}_V(m))$ in this special case.

2. IRRATIONAL ASYMPTOTIC BEHAVIOUR

In this section, we will give examples, stated in Theorems 0.2 and 0.3 of the introduction, in which the limit proved to exist in Section 1 is an irrational number. This exhibits how complicated the length $\lambda(H_m^0(R/I^n))$ can be asymptotically. In fact, we will show that the construction given by the first author in [Cu] provides an example.

Let S be a K3 surface defined over the complex field \mathbb{C} with $\text{Pic}(S) \cong \mathbb{Z}^3$. We can therefore identify $\text{Pic}(S)$ with integral points $(x, y, z) \in \mathbb{Z}^3$. Take S to be the K3 surface which has the intersection form

$$q(x, y, z) = 4x^2 - 4y^2 - 4z^2, \quad (2.1)$$

where $q(D) = D^2$ for any divisor $D \in \text{Pic}(S)$. Such a surface S exists as shown in [Cu]. It is shown there that a divisor D on S is ample if and only if it is in the interior of

$$\overline{NE}(S) = \{(x, y, z) \in \mathbb{R}^3 \mid 4x^2 - 4y^2 - 4z^2 \geq 0, x \geq 0\}.$$

Moreover, S is embedded into \mathbb{P}^3 by the divisor $H = (1, 0, 0)$. Suppose $(a, b, c) \in \mathbb{Z}^3$ is such that

$$\begin{cases} a > 0, \\ a^2 - b^2 - c^2 > 0, \\ \sqrt{b^2 + c^2} \notin \mathbb{Q}. \end{cases} \quad (2.2)$$

Since (a, b, c) is in the interior of $\overline{NE}(S)$, the divisor $A = (a, b, c)$ is ample on S . Let C be a nonsingular curve on S such that $C \sim A$. Again, C exists as shown in [Cu].

Let $R = \mathbb{C}[x_1, \dots, x_4]$ be the coordinate ring of \mathbb{P}^3 , and let I be the defining ideal of C in \mathbb{P}^3 . We will show that there exist $(a, b, c) \in \mathbb{Z}^3$ satisfying (2.2) and a curve C as above, such that

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^4} \notin \mathbb{Q}, \quad (2.3)$$

where $\mathfrak{m} = (x_1, \dots, x_4)$ is the maximal homogeneous ideal of R . Theorem 0.2 is thus an immediate consequence.

As in proved in (1.5), (1.6) and (2.1) of Theorem 1.3,

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_{\mathfrak{m}}^0(R/I^n))}{n^4} = \lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^4} + \lim_{n \rightarrow \infty} \frac{\tau(n)}{n^4},$$

where $\lim_{n \rightarrow \infty} \frac{\tau(n)}{n^4} \in \mathbb{Q}$. It remains to show that there exist $(a, b, c) \in \mathbb{Z}^3$ satisfying (2.2) and a curve C as above, such that

$$\lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^4} \notin \mathbb{Q}.$$

Let \mathcal{I} be the ideal sheaf of I on \mathbb{P}^3 . Let $\pi : X \rightarrow \mathbb{P}^3$ be the blowing up of \mathbb{P}^3 along the ideal sheaf \mathcal{I} . There exists a hyperplane H' of \mathbb{P}^3 such that $H' \cdot S = H$. Let \tilde{H} be the pull-back to X of H' , and E the exceptional divisor of the blowing up. Let $\lambda_1 = a - \sqrt{b^2 + c^2}$ and $\lambda_2 = a + \sqrt{b^2 + c^2}$. The following facts were proved in [Cu].

Lemma 2.1. *Suppose $\lambda_2 > 7$. Then,*

- (1) $h^0(S, \mathcal{O}_S(mH - nC)) = 0$ if $m < \lambda_2 n$.
- (2) $h^0(S, \mathcal{O}_S(mH - nC)) = \frac{1}{2}(mH - nC)^2 + 2$ if $m > \lambda_2 n$.
- (3) $h^1(X, \mathcal{O}_X(m\tilde{H} - nE)) = 0$ if $m > \lambda_2 n$.

Proof. (1) follows from [Cu, Remark 6]. (2) is a consequence of [Cu, Theorem 7]. (3) follows from [Cu, Theorem 9]. \square

It was pointed out in [Cu, (11)] that

$$\pi_* \mathcal{O}_X(m\tilde{H} - nE) \cong \mathcal{I}^n(m), \quad (2.4)$$

$$R^i \pi_* \mathcal{O}_X(m\tilde{H} - nE) = 0, \text{ for } i > 0. \quad (2.5)$$

Thus, we can use the cohomology groups of $\mathcal{O}_X(m\tilde{H} - nE)$ to calculate $\sigma(n)$. For convenience, we will use $H^i(X, m\tilde{H} - nE)$ and $H^i(S, mH - nC)$ to denote $H^i(X, \mathcal{O}_X(m\tilde{H} - nE))$ and $H^i(S, \mathcal{O}_S(mH - nC))$, respectively. When there is no danger of confusion, we shall further omit the space X and S in these cohomology groups. It was also shown in [Cu, (12)] that there exists the following exact sequence:

$$0 \rightarrow \mathcal{O}_X((m-4)\tilde{H} - (n-1)E) \rightarrow \mathcal{O}_X(m\tilde{H} - nE) \rightarrow \mathcal{O}_S(mH - nC) \rightarrow 0. \quad (2.6)$$

The existence of the desired example follows from the following theorem.

Theorem 2.2. *There exist $(a, b, c) \in \mathbb{Z}^3$ satisfying (2.2) and a corresponding nonsingular curve C such that, if $I \subset R$ is the defining ideal of C , then*

$$\lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^4} \notin \mathbb{Q},$$

and

$$\lim_{n \rightarrow \infty} \frac{\lambda(H_m^0(R/I^n))}{n^4} \notin \mathbb{Q}.$$

Proof. Taking the long exact sequence of cohomology groups from the exact sequence (2.6), we get

$$\begin{aligned} 0 \rightarrow H^0((m-4)\tilde{H} - (n-1)E) \rightarrow H^0(m\tilde{H} - nE) \rightarrow \\ H^0(mH - nC) \rightarrow H^1((m-4)\tilde{H} - (n-1)E). \end{aligned}$$

It follows from Lemma 2.1 that

$$H^0(m\tilde{H} - nE) \cong H^0((m-4)\tilde{H} - (n-1)E) \text{ if } m < \lambda_2 n, \quad (2.7)$$

$$H^0(m\tilde{H} - nE) \cong H^0((m-4)\tilde{H} - (n-1)E) \oplus H^0(mH - nC) \text{ if } m > \lambda_2 n. \quad (2.8)$$

Write $m = 4n + r$. Consider the following cases.

Case 1: $r < 0$. Since $\lambda_2 > 4$, we have $m < \lambda_2 n$. Thus, using (2.7) and successive induction, we get

$$H^0(m\tilde{H} - nE) = H^0(r\tilde{H}) = H^0(\mathcal{O}_{\mathbb{P}^3}(r)) = 0. \quad (2.9)$$

Case 2: $r \geq 0$. If $r > (\lambda_2 - 4)n$, i.e. $m > \lambda_2 n$, then using (2.8) and successive induction, we get $h^0(m\tilde{H} - nE) = \sum_{k=1}^n h^0((r+4k)H - kC) + 2n + h^0(\mathcal{O}_{\mathbb{P}^3}(r))$. Lemma 2.1 now gives

$$h^0(m\tilde{H} - nE) = \frac{1}{2} \sum_{k=1}^n ((r+4k)H - kC)^2 + 2n + h^0(\mathcal{O}_{\mathbb{P}^3}(r)). \quad (2.10)$$

On the other hand, if $r < (\lambda_2 - 4)n$, then put $t = \lfloor \frac{r}{\lambda_2 - 4} \rfloor$. By successive induction using both (2.7) and (2.8), we get

$$\begin{aligned} h^0(m\tilde{H} - nE) &= \sum_{k=1}^t h^0((r+4k)H - kC) + h^0(\mathcal{O}_{\mathbb{P}^3}(r)) \\ &= \frac{1}{2} \sum_{k=1}^t ((r+4k)H - kC)^2 + 2t + h^0(\mathcal{O}_{\mathbb{P}^3}(r)). \end{aligned}$$

By (2.4), we have

$$\begin{aligned}
\sigma(n) &= \sum_{m=0}^{en} h^0(m\tilde{H} - nE) \\
&= \sum_{r=-4n}^{(e-4)n} h^0((r+4n)\tilde{H} - nE) \\
&= \sum_{r=-4n}^{-1} h^0((r+4n)\tilde{H} - nE) + \sum_{r=0}^{[(\lambda_2-4)n]} h^0((r+4n)\tilde{H} - nE) \\
&\quad + \sum_{r=[(\lambda_2-4)n]+1}^{(e-4)n} h^0((r+4n)\tilde{H} - nE).
\end{aligned}$$

This together with (2.9), (2.10) and (2) gives us

$$\begin{aligned}
\sigma(n) &= \sum_{r=0}^{[(\lambda_2-4)n]} \left(\frac{1}{2} \sum_{k=1}^{\lfloor \frac{r}{\lambda_2-4} \rfloor} ((r+4k)H - kC)^2 + 2 \left\lfloor \frac{r}{\lambda_2-4} \right\rfloor + h^0(\mathcal{O}_{\mathbb{P}^3}(r)) \right) \\
&\quad + \sum_{r=[(\lambda_2-4)n]+1}^{(e-4)n} \left(\frac{1}{2} \sum_{k=1}^n ((r+4k)H - kC)^2 + 2n + h^0(\mathcal{O}_{\mathbb{P}^3}(r)) \right) \\
&= \sum_{r=0}^{[(\lambda_2-4)n]} \left(\frac{1}{2} \sum_{k=1}^{\lfloor \frac{r}{\lambda_2-4} \rfloor} ((r+4k)H - kC)^2 \right) \\
&\quad + \sum_{r=[(\lambda_2-4)n]+1}^{(e-4)n} \left(\frac{1}{2} \sum_{k=1}^n ((r+4k)H - kC)^2 \right) + \sum_{r=0}^{(e-4)n} h^0(\mathcal{O}_{\mathbb{P}^3}(r)) \\
&\quad + 2 \left(\sum_{r=0}^{[(\lambda_2-4)n]} \left\lfloor \frac{r}{\lambda_2-4} \right\rfloor + n((e-4)n - [(\lambda_2-4)n] - 1) \right).
\end{aligned}$$

Let

$$Q(s, r) = \frac{1}{2} \sum_{k=1}^s ((r+4k)H - kC)^2,$$

$$V(n) = 2 \left(\sum_{r=0}^{[(\lambda_2-4)n]} \left\lfloor \frac{r}{\lambda_2-4} \right\rfloor + n((e-4)n - [(\lambda_2-4)n] - 1) \right)$$

and

$$U(n) = \sum_{r=0}^{(e-4)n} h^0(\mathcal{O}_{\mathbb{P}^3}(r)).$$

Let

$$P(s, r) = Q(s, r) - Q(s-1, r) = \frac{1}{2} ((r+4s)H - sC)^2$$

with the convention that $Q(s, r) = 0$ for $s < 1$. For simplicity, let us also denote $\lambda = \lambda_2 - 4$. Then, we can rewrite $\sigma(n)$ as follows:

$$\sigma(n) = \sum_{r=[\lambda]+1}^{[\lambda n]} P(1, r) + \cdots + \sum_{r=[\lambda(n-1)]+1}^{[\lambda n]} P(n-1, r) + \sum_{r=[\lambda n]+1}^{(e-4)n} Q(n, r) + U(n) + V(n). \quad (2.11)$$

Let us consider one term $\sum_{r=[\lambda l]+1}^{[\lambda n]} P(l, r)$ of the sum (2.11) for some $1 \leq l \leq n-1$. From the intersection form q of (2.1) we have

$$\begin{aligned} \sum_{r=[\lambda l]+1}^{[\lambda n]} P(l, r) &= \sum_{r=[\lambda l]+1}^{[\lambda n]} 2((r + (4-a)l)^2 - (b^2 + c^2)l^2) \\ &= \sum_{r=1}^{[\lambda n]} 2((r + (4-a)l)^2 - (b^2 + c^2)l^2) - \sum_{r=1}^{[\lambda l]} 2((r + (4-a)l)^2 - (b^2 + c^2)l^2) \\ &= \left(\frac{2}{3}([\lambda n])^3 + 2(4-a)l([\lambda n])^2 + 2((4-a)^2 - b^2 - c^2)l^2([\lambda n]) \right) \\ &\quad - \left(\frac{2}{3}([\lambda l])^3 + 2(4-a)l([\lambda l])^2 + 2((4-a)^2 - b^2 - c^2)l^2([\lambda l]) \right) \\ &\quad + [\lambda n]^2 + \left(\frac{1}{3} + 2(4-a)l \right) [\lambda n] + [\lambda l]^2 + \left(\frac{1}{3} + 2(4-a)l \right) [\lambda l]. \end{aligned}$$

Note that $\lambda l - 1 < [\lambda l] < \lambda l$ for any l . Thus, we have

$$\begin{aligned} \sum_{r=[\lambda l]+1}^{[\lambda n]} P(l, r) &= \left(\frac{2}{3}(\lambda n)^3 + 2(4-a)l(\lambda n)^2 + 2((4-a)^2 - b^2 - c^2)l^2(\lambda n) \right) \\ &\quad - \left(\frac{2}{3}(\lambda l)^3 + 2(4-a)l(\lambda l)^2 + 2((4-a)^2 - b^2 - c^2)l^2(\lambda l) \right) \\ &\quad + F(n, l) \end{aligned}$$

where $F(n, l)$ is a function such that there exists a polynomial $G(n, l)$ of degree 2 with positive real coefficients satisfying $|F(n, l)| < G(n, l)$ for all $n, l \in \mathbf{N}$.

Taking the sum as l goes from 1 to $(n-1)$, we get

$$\begin{aligned} \sum_{l=1}^{n-1} \sum_{r=[\lambda l]+1}^{[\lambda n]} P(l, r) &= \left(\frac{2}{3}\lambda^3 n^4 + (4-a)\lambda^2 n^4 + \frac{2}{3}((4-a)^2 - b^2 - c^2)\lambda n^4 \right) \\ &\quad - \left(\frac{1}{6}\lambda^3 n^4 + \frac{1}{2}(4-a)\lambda^2 n^4 + \frac{1}{2}((4-a)^2 - b^2 - c^2)\lambda n^4 \right) + O(n^3). \end{aligned}$$

We also have

$$\begin{aligned} Q(n, r) &= \sum_{k=1}^n 2((r + (4-a)k)^2 - (b^2 + c^2)k^2) \\ &= 2r^2 n + 2(4-a)rn^2 + \frac{2}{3}((4-a)^2 - b^2 - c^2)n^3 + H(n, r) \end{aligned}$$

where $H(n, r)$ is a real polynomial of degree ≤ 2 in n and r . Thus,

$$\begin{aligned} \sum_{r=[\lambda n]+1}^{(e-4)n} Q(n, r) &= \sum_{r=1}^{(e-4)n} Q(n, r) - \sum_{r=1}^{[\lambda n]} Q(n, r) \\ &= \left(\frac{2}{3}(e-4)^3 n^4 + (4-a)(e-4)^2 n^4 + \frac{2}{3}((4-a)^2 - b^2 - c^2)(e-4)n^4 \right) \\ &\quad - \left(\frac{2}{3}n([\lambda n])^3 + (4-a)n^2([\lambda n])^2 + \frac{2}{3}((4-a)^2 - b^2 - c^2)n^3[\lambda n] \right) + O(n^3) \\ &= An^4 - \left(\frac{2}{3}\lambda^3 n^4 + (4-a)\lambda^2 n^4 + \frac{2}{3}((4-a)^2 - b^2 - c^2)\lambda n^4 \right) + O(n^3), \end{aligned}$$

where $A \in \mathbb{Q}$. Moreover,

$$U(n) = \sum_{r=0}^{(e-4)n} h^0(\mathcal{O}_{\mathbb{P}^3}(r)) = \sum_{r=0}^{(e-4)n} \binom{r+3}{3} = \frac{1}{24}(e-4)^4 n^4 + O(n^3) = Bn^4 + O(n^3)$$

with $B \in \mathbb{Q}$. Further, we have $V(n) = O(n^2)$. Hence,

$$\sigma(n) = (A+B)n^4 - \left(\frac{1}{6}\lambda^3 + \frac{1}{2}(4-a)^2\lambda^2 + \frac{1}{2}((4-a)^2 - b^2 - c^2)\lambda \right) n^4 + O(n^3).$$

Finally, take $a = 4, b = 3$ and $c = 2$, then clearly (a, b, c) satisfies all the requirements in (2.2) and $\lambda_2 = a + \sqrt{b^2 + c^2} > 7$. We have $\sigma(n) = (A+B)n^4 + \frac{13\sqrt{13}}{3}n^4 + O(n^3)$, where $A, B \in \mathbb{Q}$. Therefore,

$$\lim_{n \rightarrow \infty} \frac{\sigma(n)}{n^4} = A + B + \frac{13\sqrt{13}}{3} \notin \mathbb{Q}.$$

The theorem is proved. \square

We now prove Theorem 0.3. Let I and R be the ideal and ring of this section. Let $\mathfrak{m} = (x_1, \dots, x_r)$, $S = R_{\mathfrak{m}}$, $J = I_{\mathfrak{m}}$. Let E be the injective hull of \mathbb{C} .

$$\mathrm{Hom}_S(H_{\mathfrak{m}S}^0(S/J^n), E) \cong \mathrm{Ext}_S^d(S/J^n, S)$$

by local duality (c.f. Theorem 3.5.8 [BH]). Thus

$$\lambda(\mathrm{Ext}_S^d(S/J^n, S)) = \lambda(H_{\mathfrak{m}S}^0(S/J^n)) = \lambda(H_{\mathfrak{m}}^0(R/I^n))$$

for all n (c.f. Proposition 3.2.12 [BH]). Now the theorem follows from Theorem 0.2.

REFERENCES

- [BH] W. Bruns and J. Herzog. *Cohen-Macaulay rings*. Cambridge studies in advanced mathematics **39**, Cambridge University Press, Cambridge.
- [Ch] K. A. Chandler. *Regularity of the powers of an ideal*. Commun. Algebra. **25** (1997), 3773-3776.
- [CHTV] A. Conca, J. Herzog, N.V. Trung and G. Valla. *Diagonal subalgebras of bi-graded algebras and embeddings of blow-ups of projective spaces*. American Journal of Math. **119** (1997), 859-901.
- [Cu] S.D. Cutkosky. *Irrational asymptotic behaviour of Castelnuovo-Mumford regularity*. J. reine angew. Math. **522** (2000), 93-103.
- [CEL] S.D. Cutkosky, L. Ein, and R. Lazarsfeld. *Positivity and complexity of ideal sheaves*. Math. Ann. **321** (2001), 213-234.
- [CHT] Cutkosky, S.D., Herzog, J. and Trung, N.V. (1999). *Asymptotic behaviour of the Castelnuovo-Mumford regularity*. Compositio Math. **118**, 243-261.
- [Fu] T. Fujita. *Approximating Zariski decomposition of big line bundles*. Kodai Math. J. **17** (1994), 1-3.

- [F2] T. Fujita. *Semipositive line bundles*. Journal of the Faculty of Science, The University of Tokyo **30** (1983), 353-378.
- [GGP] A.V. Geramita, A. Gimigliano, and Y. Pitteloud. *Graded Betti numbers of some embedded rational n -folds*. Math. Ann. **301** (1995), 363-380.
- [HaT] H. Tàì Hà and N.V. Trung. *Arithmetic Cohen-Macaulayness of blow-ups*. Preprint.
- [H] R. Hartshorne. *Algebraic Geometry*. GTM 52 Springer Verlag (1977) Berlin Heidelberg.
- [HoH] L.T. Hoa and E. Hyry. *On local cohomology and Hilbert function of powers of ideals*. Preprint.
- [HgT] N.D. Hoang and N.V. Trung. *Hilbert polynomials of non-standard bigraded algebras*. Preprint. [aXiv:math.AC/0211181](https://arxiv.org/abs/math/0211181).
- [I] S. Iitaka, *Algebraic Geometry*. GTM 76, Springer Verlag, New York, Heidelberg, Berlin.
- [Ki] D. Kirby. *Hilbert functions and the extension functor*. Math. Proc. Camb. Phil. Soc. **105** (1989), no. 3, 441-446.
- [Ko1] V. Kodiyalam. *Homological invariants of powers of an ideal*. Proc. Amer. Math. Soc. **118** (1993), no. 3, 757-764.
- [Ko2] V. Kodiyalam. *Asymptotic behaviour of Castelnuovo-Mumford regularity*. Proc. Amer. Math. Soc. **128** (2000), no. 2, 407-411.
- [La] R. Lazarsfeld. *Positivity in Algebraic Geometry*. Book to be published.
- [M] S. McAdam. *Asymptotic prime divisors and analytic spreads*. Proc. AMS **80** (1980), 555-559.
- [STV] A. Simis, N.V. Trung and G. Valla. *The diagonal subalgebra of a blow-up algebra*. J. Pure Appl. Algebra. **125** (1998), no. 1-3, 305-328.
- [S] I. Swanson, *Powers of ideals: Primary decompositions, Artin-Rees Lemma and Regularity*. Math. Annalen **307** (1997), 299-313.
- [Th] E. Theodorescu. *Derived functors and Hilbert polynomials*. Math. Proc. Camb. Phil. Soc. **132** (2002), 75-88.
- [Th2] E. Theodorescu. *Derived functors and Hilbert polynomials*, University of Kansas thesis, (2002).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65201, USA

E-mail address: cutkoskys@missouri.edu, tai@math.missouri.edu, srinivasanh@missouri.edu, theodore@math.missouri.edu