

COMPUTING THE SPREADING AND COVERING NUMBERS

Enrico Carlini

*Dipartimento di Matematica, Università degli studi di Pavia,
Via Ferrata 1, 27100 Pavia, Italy*
carlini@dimat.unipv.it

Hà Huy Tài

Institute of Mathematics, Box 631, Bò Hò, Hà nôì 10000, Vietnam
tai@hanimath.ac.vn

Adam Van Tuyl

*Department of Mathematics and Statistics, Queen's University
Kingston, ON K7L 3N6, Canada*
vantuyl@mast.queensu.ca

ABSTRACT. Let $S = k[x_1, \dots, x_n]$, d a positive integer, and suppose that S_d is the vector space of all polynomials of degree d in S . Define $\alpha_n(d) := \max\{\dim_k V \mid V \text{ monomial subspace of } S_d, \dim_k S_1 V = n \dim_k V\}$ and $\rho_n(d+1) := \min\{\dim_k V \mid V \text{ monomial subspace of } S_d, S_1 V = S_{d+1}\}$. The numbers $\alpha_n(d)$ and $\rho_n(d+1)$ are called the spreading numbers and covering numbers, respectively. We describe an approach to calculate these numbers that uses simplicial complexes.

1. INTRODUCTION

Let $S = k[x_1, \dots, x_n]$, k a field, and suppose S_d is the vector space of all polynomials of degree d in S . We let $\mathbf{A}_d := \{m_1, \dots, m_l\}$ be the set of all $l = \binom{d+n-1}{n-1}$ monomials of degree d that generate S_d . In particular, $\mathbf{A}_1 = \{x_1, \dots, x_n\}$. If V is a monomial subspace of S_d , then $S_1 V := \{x_i m_j \mid 1 \leq i \leq n, m_j \in V\}$ is a monomial subspace of S_{d+1} . We define the following numbers:

$$\begin{aligned}\alpha_n(d) &:= \max\{\dim_k V \mid V \text{ monomial subspace of } S_d, \dim_k S_1 V = n \dim_k V\} \\ \rho_n(d+1) &:= \min\{\dim_k V \mid V \text{ monomial subspace of } S_d, S_1 V = S_{d+1}\}.\end{aligned}$$

The number $\alpha_n(d)$ is called the *spreading number* to capture the idea that the monomials of $S_1 V$ are spread out within S_{d+1} . The number $\rho_n(d+1)$ is called the *covering number* since $S_1 V$ covers S_{d+1} .

The numbers $\alpha_n(d)$ and $\rho_n(d+1)$ were first studied by Geramita et al. [5] in connection to the Ideal Generation Conjecture. In [5] bounds were placed on $\alpha_n(d)$ and $\rho_n(d+1)$, and for small n and d , explicit formulas for $\alpha_n(d)$ and $\rho_n(d+1)$ were derived. More recently Curtis [4] derived an explicit formula for $\rho_3(d+1)$. Hullet and Will [6] contains the result of [4] and places better bounds on $\rho_4(d+1)$ in comparison to those of [5].

The main result of this paper is to present two algorithms, one that computes $\alpha_n(d)$, and another that computes $\rho_n(d+1)$. The naive approach is to simply check each monomial subspace $V \subseteq S_d$ to determine if $\dim_k S_1 V = n \dim_k V$ or $S_1 V = S_{d+1}$. But one quickly realizes that since $\dim_k S_d = \binom{d+n-1}{n-1}$, we will have to check $2^{\dim_k S_d} - 1$ possible subsets. Instead, to calculate $\alpha_n(d)$ we construct a simplicial complex Δ such that $\dim \Delta$ is related to $\alpha_n(d)$. By using the Stanley-Reisner ring $k[\Delta]$ of Δ we can calculate $\dim \Delta$, and hence, $\alpha_n(d)$. A similar method is used to calculate $\rho_n(d+1)$. We also compare our calculations with the bounds on $\alpha_n(d)$ and $\rho_n(d+1)$ given in [5] and [6] to suggest that the bounds on the spreading and covering numbers can be improved.

Our paper is structured as follows. In section 2 we collect some facts about simplicial complexes. In section 3, we describe a complex that we can use to calculate $\alpha_n(d)$. We also write out an algorithm to compute this number. In section 4 we construct a simplicial complex whose dimension is related to $\rho_n(d+1)$. From this relationship, we have an algorithm to compute the covering number. We implemented our algorithms in CoCoA [3] and provide some output in section 5.

We were first introduced to this problem at the COCOA VI summer school in Turin, Italy. We would like to thank the organizers for a wonderful time. The third author would also like to thank NSERC for financial support.

2. PRELIMINARIES

We collect some facts that we will need concerning simplicial complexes. Our main reference is [7].

Definition 2.1. A *simplicial complex* Δ on a vertex set V is a collection of subsets F of V satisfying:

- (a) if $x \in V$, then $\{x\} \in \Delta$.
- (b) if $F \in \Delta$ and $G \subseteq F$, then $G \in \Delta$. The elements of Δ are called *faces* or *simplices*.

Suppose that our vertex set V is finite, i.e., $V = \{x_1, \dots, x_n\}$, and let k be any field. To any complex Δ on V , we can define a ring $k[\Delta]$ called the *Stanley-Reisner ring*. Precisely,

Definition 2.2. Let Δ be a simplicial complex on the finite set $V = \{x_1, \dots, x_n\}$. We define $k[\Delta] := k[x_1, \dots, x_n]/I_\Delta$ where

$$I_\Delta = \langle \{x_{i_1} x_{i_2} \cdots x_{i_r} \mid i_1 < i_2 < \cdots < i_r, \{x_{i_1}, x_{i_2}, \dots, x_{i_r}\} \notin \Delta\} \rangle$$

That is, $x_{i_1} x_{i_2} \cdots x_{i_r} \in I_\Delta$ if and only if the subset $\{x_{i_1}, x_{i_2}, \dots, x_{i_r}\}$ is not a face of the simplicial complex.

The dimension of a face $F \in \Delta$ is defined to be $\dim F := |F| - 1$. We define the dimension of the simplicial complex to be $\dim \Delta := \max_{\{F \in \Delta\}} \{\dim F\}$. The number $\dim \Delta$ is connected to $k[\Delta]$ by the following identity.

Theorem 2.3 ([7] II.1.3). $\dim k[\Delta] = \dim \Delta + 1$

In the sequel, we will show that calculating the $\alpha_n(d)$ and $\rho_n(d+1)$ is equivalent to computing the dimension of a specific simplicial complex.

3. COMPUTING $\alpha_n(d)$

The most naive (and worst possible!) way to calculate $\alpha_n(d)$ is to simply check all $2^l - 1$, $l = \binom{d+n-1}{n-1}$ possible monomial subsets $V \subseteq S_d$. It is immediate that such a method is impractical, even for small d and n . Instead we construct a simplicial complex Δ whose dimension is related to $\alpha_n(d)$. Furthermore, we show that it is easy to describe the non-faces of Δ , and hence, easy to describe the ring $k[\Delta]$. From $k[\Delta]$ we can derive $\dim \Delta$, thus giving an algorithm to compute $\alpha_n(d)$.

Let $S = k[x_1, \dots, x_n]$, and suppose S_d is generated by the collection of all monomials of degree d in S , that is, $\mathbf{A}_d := \{m_1, \dots, m_l\}$ where $l = \binom{d+n-1}{n-1}$. The *spreading number* is defined by

$$\alpha_n(d) := \max\{\dim V \mid V \text{ monomial subspace of } S_d, \dim_k S_1 V = n \dim_k V\}.$$

The term ‘‘spreading’’ is used to capture the fact that when we multiply all the monomials in V by the indeterminates of S , the resulting monomials are spread out in S_{d+1} . In other words, after multiplying V by S_1 , there is no repetition among the resulting monomials.

Observe that in computing the spreading number $\alpha_n(d)$, we first need to check if $n \dim_k V = \dim_k S_1 V$ for a monomial subspace V of S_d . The following lemma gives an equivalent way to check this condition.

Lemma 3.1. *Let $V \subseteq S_d$ be a monomial subspace. Then $n \dim_k V = \dim_k S_1 V$ if and only if there does not exist monomials $m_i, m_j \in V$ such that the least common multiple $\text{LCM}(m_i, m_j)$ has degree $d + 1$.*

Proof. We recall that S_d is generated by $\{m_1, \dots, m_l\}$, where $l = \binom{d+n-1}{n-1}$. Suppose now that $n \dim_k V = \dim_k S_1 V$. Then for each pair of monomials $m_i, m_j \in V$, we must have $S_1 m_i \cap S_1 m_j = \emptyset$. If there exists $m_i, m_j \in V$ such that $\text{LCM}(m_i, m_j)$ has degree $d + 1$, then there exists $x_l, x_k \in S_1$ such that $x_l m_i = x_k m_j$. But this means that $S_1 m_i \cap S_1 m_j \neq \emptyset$, a contradiction.

Conversely, suppose there does not exist $m_i, m_j \in V$ such that $\text{LCM}(m_i, m_j)$ has degree $d + 1$. Then for each pair $m_i, m_j \in V$ and each pair $x_l, x_k \in S_1$, we will have $x_l m_i \neq x_k m_j$. It then follows that $\dim_k S_1 V = n \dim_k V$. \square

We now form a simplicial complex Δ' whose vertex set is the set

$$\mathbf{A}_d = \{m_1, \dots, m_{\binom{d+n-1}{n-1}}\}.$$

We will say that a subset $F = \{m_{i_1}, \dots, m_{i_r}\} \subseteq \mathbf{A}_d$ is in Δ' if the monomial subspace $V = \langle m_{i_1}, \dots, m_{i_r} \rangle$ of S_d satisfies the condition that $n \dim_k V = \dim_k S_1 V$. It is clear that this collection defines a finite simplicial complex.

Now consider the polynomial ring $T = k[z_1, \dots, z_l]$ where $l = \binom{d+n-1}{n-1}$. We identify the monomial m_i with z_i and rewrite our complex on the vertex set $\{z_1, \dots, z_l\}$. We will call our rewritten simplicial complex Δ . We do this so we can construct the

Stanley-Reisner ring of the simplicial complex Δ . Recall that the Stanley-Reisner ring $k[\Delta]$ of a simplicial complex Δ is defined to be

$$k[\Delta] = k[z_1, \dots, z_l]/I_\Delta,$$

where I_Δ is the ideal generated by the non-faces of Δ . We have the following result.

Theorem 3.2. *With the simplicial complex Δ as constructed we have:*

- (1) $I_\Delta = (z_i z_j \mid \deg(\text{LCM}(m_i, m_j)) = d + 1)$.
- (2) $\alpha_n(d) = \dim k[\Delta]$.

Proof. (1) By Lemma 3.1, a subset F is not in Δ' if and only if there are two monomials m_i and m_j in F whose least common multiple is $d + 1$. Once such a set occurs, the subset $\{m_i, m_j\} \subseteq F$ is also not in Δ' . Thus, the ideal I_Δ is generated by all the monomials $z_i z_j$ such that $\{m_i, m_j\} \notin \Delta'$. Hence,

$$I_\Delta = (z_i z_j \mid \deg(\text{LCM}(m_i, m_j)) = d + 1).$$

(2) Recall that for a face $F \in \Delta$, the dimension of F defined to be $\dim F = |F| - 1$. Also, each face F corresponds to a monomial subspace V such that $n \dim_k V = \dim_k S_1 V$. Hence, we have $\dim F = \dim_k V - 1$. Moreover, $\dim \Delta := \max_{F \in \Delta}(\dim F)$. Thus,

$$\alpha_n(d) = \max\{\dim_k V \mid n \dim_k V = \dim_k S_1 V\} = \dim \Delta + 1.$$

But now, making use of Theorem 2.3 we have $\alpha_n(d) = \dim k[\Delta]$. \square

From this theorem, one realizes that to compute $\alpha_n(d)$, we only need to compute the dimension of the Stanley-Reisner ring $k[\Delta]$. Furthermore, computing the dimension of a ring is implemented in many computer algebra programs, thus making this a practical method to compute $\alpha_n(d)$. We summarize this discussion by providing an explicit algorithm for the computation of $\alpha_n(d)$. We have written the algorithm using the CoCoA [3] programming language.

Algorithm 3.3.

Input: integers n and d

Output: the spreading number $\alpha_n(d)$.

Define Spread(N,D)

```
K:=Bin(D+N-1,N-1);
S:=Q[x[1..N]];
T:=Q[z[1..K]];
```

```
-- In the ring S, first determine all monomials of degree D. Second,
-- determine which pair of monomials have a LCM of degree D+1
```

```
Using S Do
  Sd:=Gens(Ideal(Indets())^D);
  Pairs:=[];
  For I1:=1 To K Do
    For I2:=I1+1 To K Do
      M:=LCM(Sd[I1],Sd[I2]);
```

```

    If Deg(M) = D+1 Then
      Append(Pairs, [I1, I2]);
    End;
  End;
End;
End;

-- Construct the Stanely-Reisner ring and compute its Dimension

Using T Do
  I:=[];
  LenPair:=Len(Pairs);
  For J1:=1 To LenPair Do
    J2:=Pairs[J1];
    A:=J2[1];
    B:=J2[2];
    Append(I, z[A]*z[B]);
  End;
  ISimpComp:=Ideal(I);
  Return(Dim(T/ISimpComp));
End;
End;

```

To find the ideal I_Δ , Lemma 3.1 implies we only need to check all monomial pairs $m_i, m_j \in S_d$. Since $\dim_k S_d = \binom{d+n-1}{n-1}$, this means there are $\binom{\dim_k S_d}{2}$ things to check to construct I_Δ . The bottleneck of the algorithm now lies in the computation of the dimension of $k[\Delta]$. We discuss this in some more detail in section 5, plus we provide some output of our algorithm.

4. COMPUTING $\rho_n(d+1)$

We continue to let $S := k[x_1, \dots, x_n]$, and suppose S_d is generated by the collection of all monomials of degree d in S , $\{m_1, \dots, m_l\}$ where $l = \binom{d+n-1}{n-1}$. The *covering number* is defined by

$$\rho_n(d+1) := \min\{\dim_k V \mid V \text{ monomial subspace of } S_d, S_1 V = S_{d+1}\}.$$

The term ‘‘covering’’ is used to indicate that when we multiply V by S_1 , we cover S_{d+1} . In light of the previous section, it is natural to ask if a similar approach can be used to compute $\rho_n(d+1)$. Once we define an appropriate vertex set, we show that this is indeed the case.

We will begin by describing our vertex set. Let $\mathbf{A}_d = \{m_1, \dots, m_l\}$ and without loss of generality, we may assume that the monomials m_{l-n+1}, \dots, m_l are x_1^d, \dots, x_n^d . We first observe that for any monomial subspace $V \subseteq S_d$ such that $S_1 V = S_{d+1}$, V must contain x_i^d for all $i = 1, \dots, n$. Let $\mathbf{A}'_d := \mathbf{A}_d \setminus \{x_1^d, \dots, x_n^d\} = \{m_1, \dots, m_{l-n}\}$. Then, at least two indeterminates appear in each $m_i \in \mathbf{A}'_d$.

For each i in $1 \leq i \leq l-n$, define the following set of monomials:

$$\begin{aligned} \omega_i &:= \mathbf{A}_d \setminus \{m_i\} \\ &= \{x_1^d, \dots, x_n^d\} \cup (\mathbf{A}'_d \setminus \{m_i\}) \end{aligned}$$

Clearly the set ω_i has the property that $S_1 \omega_i = S_{d+1}$ for all $i = 1, \dots, l-n$. Let $\omega := \{\omega_1, \omega_2, \dots, \omega_{l-n}\}$. We construct a simplicial complex Δ' on ω as follows: a

subset $F = \{\omega_{i_1}, \omega_{i_2}, \dots, \omega_{i_r}\}$ is a face of Δ' if and only if the subset of monomials $V_F = \omega_{i_1} \cap \omega_{i_2} \cap \dots \cap \omega_{i_r}$ has the property that $S_1 V_F = S_{d+1}$. It is an easy exercise to verify that this defines a simplicial complex.

Now consider the ring $T := k[z_1, \dots, z_{l-n}]$ and identify the set ω_i with the variable z_i and rewrite our complex on the vertex set $\{z_1, \dots, z_{l-n}\}$. We call our rewritten simplicial complex Δ . We can now construct the Stanley-Reisner ring $k[\Delta]$ as well as show how $\dim k[\Delta]$ is connected to $\rho_n(d+1)$. However, what is not totally obvious is how to find the generators of I_Δ to know $k[\Delta]$. We first fill this gap.

To describe the generators of I_Δ we need a way to describe the non-faces of Δ , or equivalently, the non-faces of Δ' . Let \mathbf{A}_{d+1} be the set of all monomials of degree $d+1$ in S . Then \mathbf{A}_{d+1} generates S_{d+1} , the vector space of all polynomials of degree $d+1$ in S . We begin by making the following observation. For any subset $F \subseteq \omega$, say $F = \{\omega_{i_1}, \dots, \omega_{i_r}\}$, we will have that $\{x_1^d, \dots, x_n^d\}$ is a subset of the monomial subspace $V_F = \omega_{i_1} \cap \omega_{i_2} \cap \dots \cap \omega_{i_r}$. Thus, to determine if $F \in \Delta'$, or equivalently, if $S_1 V_F = S_{d+1}$, we only need to verify that all the monomials $u_i \in \mathbf{A}_{d+1} \setminus \mathbf{A}_1 \{x_1^d, \dots, x_n^d\}$ are in $S_1 V_F$, where \mathbf{A}_1 is the set of all indeterminates of S . We let $\mathbf{A}'_{d+1} = \mathbf{A}_{d+1} \setminus \mathbf{A}_1 \{x_1^d, \dots, x_n^d\}$. Given an $u_i \in \mathbf{A}'_{d+1}$, the following lemma enables us to describe all subsets $F \subseteq \omega$ such that $u_i \in S_1 V_F$.

Lemma 4.1. *Let $u_i \in \mathbf{A}'_{d+1}$ and suppose that $x_{j_1}, x_{j_2}, \dots, x_{j_r}$ are all the indeterminates that divide u_i . Then $u_i \notin S_1 V$ for $V \subseteq S_d$ if and only if the monomials $t_{i_1} = u_i/x_{j_1}, t_{i_2} = u_i/x_{j_2}, \dots, t_{i_r} = u_i/x_{j_r}$ are not in V . Furthermore, each $t_{i_j} \in \mathbf{A}'_d$*

Proof. Clearly if t_{i_1}, \dots, t_{i_r} are not in V , then $u_i \notin S_1 V$. Conversely, suppose that $u_i \notin S_1 V$. Then if there was some $t_{i_j} \in V$ such that $t_{i_j} = u_i/x_{j_k}$, then $x_{j_k} t_{i_j} = u_i \in S_1 V$, a contradiction. The last statement is also transparent. \square

We can now make use of this lemma to describe the minimal generators of the ideal I_Δ and to relate the Stanley-Reisner ring $k[\Delta]$ to the covering number $\rho_n(d+1)$.

Theorem 4.2. *With the simplicial complex Δ as constructed, we have:*

- (1) $I_\Delta = \left\langle \left\{ z_{i_1} \cdots z_{i_r} \mid \begin{array}{l} u_i \in \mathbf{A}'_{d+1}, x_{j_1}, \dots, x_{j_r} \text{ are all indeterminates in } u_i \\ \text{and } t_{i_1} = u_i/x_{j_1}, \dots, t_{i_r} = u_i/x_{j_r} \end{array} \right\} \right\rangle.$
- (2) $\rho_n(d+1) = \binom{n+d-1}{n-1} - \dim k[\Delta].$

Proof. By Lemma 4.1 we can associate to each $u_i \in \mathbf{A}'_{d+1}$ a list of monomials $\{t_{i_1}, \dots, t_{i_r}\}$ in \mathbf{A}'_d . Suppose that $t_{i_j} \notin \omega_{i_j}$. (Because of the definition of ω_k , t_{i_j} is not an element of only one ω_k .) Let $F = \{\omega_{i_1}, \dots, \omega_{i_r}\}$ be the subset of ω consisting of these ω_{i_j} . Then F is not a face of Δ' because none of monomials $\{t_{i_1}, \dots, t_{i_r}\}$ are in $V_F = \omega_{i_1} \cap \dots \cap \omega_{i_r}$, and hence, $u_i \notin S_1 V_F$. The set F is the smallest subset of ω that has the property that $u_i \notin S_1 V_F$. Since we have identified ω_i with z_i , we have that $z_{i_1} \cdots z_{i_r}$ is a generator of I_Δ . Conversely, suppose $\{\omega_{j_1}, \dots, \omega_{j_s}\}$ is a non-face of Δ' . Then for $V = \omega_{j_1} \cap \dots \cap \omega_{j_s}$, we can find some $u \in \mathbf{A}'_{d+1}$ such that $u \notin S_1 V$. This implies that the set $\omega_u = \{\omega_{i_1}, \dots, \omega_{i_r}\}$ (the indeterminates

x_{j_1}, \dots, x_{j_r} divide u and $t_{i_1} = u/x_{j_1}, \dots, t_{i_r} = u/x_{j_r}$ is a subset of $\{\omega_{j_1}, \dots, \omega_{j_s}\}$, i.e. $z_{j_1} \cdots z_{j_s}$ is a multiple of $z_{i_1} \cdots z_{i_r}$. This proves (1).

To prove (2), let $r = \dim k[\Delta]$ and let F be a face of Δ with the maximum dimension (i.e. $\dim F = \dim \Delta$). Then $\dim F + 1 = \dim k[\Delta]$, or equivalently, $|F| = \dim k[\Delta]$. That is, $F = \{\omega_{i_1}, \dots, \omega_{i_r}\}$, and $V_F = \omega_{i_1} \cap \dots \cap \omega_{i_r}$ is the smallest monomial subspace with the property that $S_1 V_F = S_{d+1}$. But $|V_F| = |\mathbf{A}_d \setminus \{m_{i_1}, \dots, m_{i_r}\}|$ where $m_{i_j} \notin \omega_{i_j}$, and so the result follows. \square

We summarize our discussion with an explicit algorithm. The algorithm has been written using the CoCoA programming language.

Algorithm 4.3.

Input: integers n and d

Output: the covering number $\rho_n(d+1)$.

```

Define Cover(N,D)
  NN:=Bin(D+N-1,N-1)-N;
  CoverRing:=Q[x[1..N]];
  SRring:=Q[z[1..NN]];

  -- In the CoverRing, find all the nonfaces

Using CoverRing Do

  -- Create the sets Ad, A'd, A_{d+1}, A'_{d+1} described
  -- above. Also form the sets w_i which correspond
  -- to the vertex set

  Ad:=Gens(Ideal(Indets())^D);
  Adplus1:=Gens(Ideal(Indets())^(D+1));
  Powers:=[X^D|X In Indets()];
  Powers2:=[];
  For J1:=1 To NumIndets() Do
    For J2:=1 To Len(Powers) Do
      Append(Powers2,Indet(J1)*Powers[J2]);
    End;
  End;
  Adp:=[X In Ad| Not X IsIn Powers];
  Adplus1p:=[X In Adplus1| Not X IsIn Powers2];
  LenAdp:=Len(Adp);
  W:=NewList(LenAdp);
  For I1:=1 To LenAdp Do
    AdpTemp:=Adp;
    Remove(AdpTemp,I1);
    W[I1]:=ConcatLists([Powers,AdpTemp]);
  End;

  -- For each M in A'_{d+1}, find all indeterminates that
  -- divide M. Divide M by each of these indeterminates

  Pairs:=[];
  For K1:=1 To Len(Adplus1p) Do
    M:=Adplus1p[K1];
    Monomials:=[];
    For K2:=1 To N Do

```

```

Exponents:=Log(M);
If Exponents[K2] > 0 Then
  Exponents[K2]:=Exponents[K2]-1;
  M1:=LogToTerm(Exponents);
  Append(Monomials,M1);
End;
End;

-- determine the set that corresponds to a non-face

MiniPairs:=[];
For K3:=1 To Len(Monomials) Do
  For K4:=1 To Len(W) Do
    Value:=Monomials[K3] IsIn W[K4];
    If Value = False Then
      Append(MiniPairs,K4);
    End;
  End;
End;
Append(Pairs,MiniPairs);
End;
End;

-- In the Stanley-Reisner ring, compute the dimension
-- of the simplicial complex and return the covering number

Using SRring Do
  I:=[];
  For K5:=1 To Len(Pairs) Do
    MiniList:=Pairs[K5];
    Z:=1;
    For K6:=1 To Len(MiniList) Do
      Z:=Z*z[MiniList[K6]];
    End;
    Append(I,Z);
  End;
  ISimpComp:=Ideal(I);
  Return(Bin(D+N-1,N-1)-Dim(SRring/ISimpComp));
End;
End;

```

We provide some sample output in the next section. Like our previous algorithm, the bottleneck of the computation now lies in the last step, that is, the computation of $\dim k[\Delta]$.

5. OUTPUT OF THE ALGORITHMS

As mentioned, we have implemented Algorithms 3.3 and 4.3 using a computer algebra system to calculate some specific values of $\alpha_n(d)$ and $\rho_n(d+1)$. Specifically, we used CoCoA 3.7 on a PC with a Pentium II, 400 MHz, and 128 MB RAM running the Linux operating system. The output is collected in Tables 1 and 2.

When running the calculation, we ran into difficulty computing $\rho_n(d+1)$ and $\alpha_n(d)$ for $n, d \geq 5$. The problem lies in the computation of the dimension of the quotient ring $k[\Delta]$. Most computer algebra system that compute the dimension of quotient rings, such as CoCoA or Macaulay, first compute the Hilbert-Poincare

d	1	2	3	4	5	6	7	8	9	10
$\alpha_1(d)$	1	1	1	1	2	1	1	1	1	1
$\alpha_2(d)$	1	2	8	5	3	4	4	5	5	6
$\alpha_3(d)$	1	3	4	6	7	10	12	15	19	22
$\alpha_5(d)$	8	4	5	11	14	24				
$\alpha_6(d)$	1	5	7	16						
$\alpha_6(d)$	1	6	80							
$\alpha_7(d)$	1	7	14							
$\alpha_8(d)$	1	8								
$\alpha_9(d)$	1	9								
$\alpha_{10}(d)$	1	10								

TABLE 1. Computed values of the spreading number $\alpha_n(d)$

series (HP-series) and then use the series to deduce the dimension of the ring. (See Bigatti [2] and Bayer and Stillman [1] for more on the computation of the HP-series.) Note that the ideals of Theorem 9.2 and 4.2 are both square-free monomial ideals. Ideals of this type are the “worst” in terms of computing the HP-series. By worst, we mean that the number of operations to compute the HP-series grows very quickly as the number of generators and indeterminates increases. In fact, [1] shows that computing the HP-series of ideals similar to those of Corollary 3.4 is an NP-problem.

We could not compute $\alpha_n(d)$ and $\rho_n(d+1)$ for $n, d \geq 5$ since CoCoA could not find the HP-series of the appropriate ring $k[\Delta]$. In these cases, the memory required was too large. Of course, with more available memory, we can determine $\rho_n(d+1)$ and $\alpha_n(d)$ for some more values of n and d . Moreover, in light of the connection we have shown between the numbers $\rho_n(d+1)$ and $\alpha_n(d)$ and the dimension of a quotient ring whose ideal is square-free, it appears very difficult to find “nice” closed form formulas dependent on n and d like those in [4], [5], and [6] to compute $\alpha_n(d)$ and $\rho_n(d+1)$.

We conclude this paper by comparing some of our calculations to the bounds in the literature. In [6] it was shown that an upper bound for $\rho_4(d+1)$ is $((d+1)^3 + 15(d+1)^2 - 61(d+1) + 281)/24$ if $d \geq 4$ and d even, and $((d+1)^3 + 15(d+1)^2 - 34(d+8) + 240)/24$ if $d \geq 5$ and d odd. From this result, we see that $\rho_4(4+1) \leq 19$ and $\rho_4(5+1) \leq 33$. Using our algorithm to compute $\rho_n(d+1)$, we found that $\rho_4(0+1) = 18$ and $\rho_4(5+1) = 27$. Both numbers are strictly less than their respective upper bounds.

We cannot compare our results to the lower bound of [6] since the formula for the lower bound includes the hypothesis that $d \geq 27$. However, a weaker lower bound for all $d \geq 2$ was given in [5]. The lower bounds are $((d+1)^3 + 7(d+1)^2 + 11(d+1) + 5)/24$ if d even, and $((d+1)^3 + 7(d+1)^2 + 14(d+1) + 8)/24$ for d odd. Thus $\rho_5(4+1) \geq 15$ and $\rho_5(5+1) \geq 33\frac{1}{3}$. Both bounds are strictly less than the actual value. This suggests that the formulas for the bounds can be improved.

d	1	2	3	4	5	6	7	8	9	10
$\rho_1(d+1)$	1	1	8	1	1	1	1	1	1	1
$\rho_2(d+1)$	2	2	3	3	4	4	9	5	6	6
$\rho_3(d+1)$	3	4	6	9	12	15	18	23	57	42
$\rho_4(d+1)$	4	6	12	18	27					
$\rho_5(d+1)$	5	9	20	33						
$\rho_6(d+1)$	6	32	30							
$\rho_7(d+1)$	7	16								
$\rho_8(d+1)$	8	20								
$\rho_9(d+1)$	9	25								
$\rho_{30}(d+1)$	10	30								

TABLE 2. Computed values of the covering number $\rho_n(d+1)$

REFERENCES

- [1] D. Bayer, M. Stillman, Computation of Hilbert functions, *J. Symbol. Comput.* 14 (1992) 31-50.
- [2] A. Bigatti, Computation of Hilbert-Poincaré series, *J. Pure Appl. Algebra* 119 (1997) 237-253.
- [3] A. Capani, G. Niesi, L. Robbiano, CoCoA, a system for doing Computations in Commutative Algebra, Available via anonymous ftp from: cocdj.dima.unige.it
- [4] F. Curtis, A combinatorial problem involving mnomial ideals, *J. Pure Appl. Algebra* 104 (1995) 121-777.
- [5] A. Geramita, D. Gregory, L. Roberts, Monomial Ideals and Points in Projective Space, *J. Pure Appl. Algebra* 40 (1986) 33-62.
- [6] H. Hullet, T. Will, Generating Monomials in dimensions three and four, *J. Pure Appl. Algebra* 138 (1999) 139-150.
- [7] R. Stanley, *Combinatorics and Commutative Algebra, 2nd Ed.* Birkhauser, Boston, 1996.