

PROOFS OF THREE GEODE CONJECTURES

TEWODROS AMDEBERHAN AND DORON ZEILBERGER

ABSTRACT. In the May 2025 issue of the Amer. Math. Monthly, Norman J. Wildberger and Dean Rubine introduced a new kind of multi-indexed numbers, that they call ‘Geode numbers’, obtained from the Hyper-Catalan numbers. They posed three intriguing conjectures about them, that are proved in this note.

1. INTRODUCTION

In a recent captivating Monthly article [4], by Norman J. Wildberger and Dean Rubine, the authors utilize a generating series to solve the general univariate polynomial equation. They also explored a “curious factorization” of this hyper-Catalan generating series, and in the penultimate section, they made three conjectures about this algebraic object that they termed the *Geode array*.

In this note, we prove these three conjectures. At least as interesting as the actual statements of the conjectures (now theorems) is *how we proved them*, using several important *tools of the trade*.

The first tool is the *multinomial theorem*

$$(1.1) \quad (x_1 + \cdots + x_r)^n = \sum_{\substack{m_1, \dots, m_r \geq 0 \\ m_1 + \cdots + m_r = n}} \binom{n}{m_1, \dots, m_r} x_1^{m_1} \cdots x_r^{m_r}.$$

The second tool is *constant-term extraction*, the third is *Wilf-Zeilberger (WZ) algorithmic proof theory* [5] and the last-but-not-least tool is *Lagrange Inversion* [6] that states that: *if $u(t)$ and $\Phi(t)$ are formal power series starting at t^1 and t^0 , respectively, then $u(t) = t\Phi(u(t))$ implies*

$$(1.2) \quad [t^n]u(t) = \frac{1}{n}[z^{n-1}]\Phi(z)^n.$$

Here $[z^n]F(z)$ means the coefficient of z^n in the Laurent expansion of $F(z)$. We shall use the notation $\mathbf{CT}_z F(z)$ for the constant-term of $F(z)$.

We now bring in the relevant notation adopted in [4] with a caveat that indices are shifted slightly. Consider the equation $0 = 1 - \alpha + \sum_{k \geq 1} t_k \alpha^{k+1}$

Date: June 25, 2025.

2000 Mathematics Subject Classification. Primary, Secondary.

and denote its series solution by $\alpha = \mathbf{S}[t_1, t_2, \dots]$. Letting $\mathbf{S}_1 = t_1 + t_2 + \dots$, Wildberger-Rubine proved [4, Theorem 12] the existence of a (remarkable!) factorization $\mathbf{S} - 1 = \mathbf{S}_1 \mathbf{G}$ and the factor $\mathbf{G}[t_1, t_2, \dots]$ (that they dubbed the *Geode series*). Furthermore, we opt to use $G[m_1, m_2, \dots]$ for the coefficient of $t_1^{m_1} t_2^{m_2} \dots$ in the polyseries $\mathbf{G}[t_1, t_2, \dots]$. We are now ready to state and prove the three conjectures from [4, p. 399]. For the sake of clarity, let's describe the first of these in some detail.

Suppose we are solving the polynomial equation $0 = 1 - \alpha + t_1 \alpha^2 + t_2 \alpha^3$ through the formal power series

$$\alpha = \mathbf{S}[t_1, t_2] = \sum_{m_1, m_2 \geq 0} C[m_1, m_2] t_1^{m_1} t_2^{m_2}.$$

Consequently, the corresponding Geode series becomes $\mathbf{G}[t_1, t_2] = \frac{\mathbf{S}[t_1, t_2] - 1}{t_1 + t_2}$. We follow closely [6] to engage the Lagrange Inversion in the extraction of the coefficients $C[m_1, m_2]$ satisfying $n = m_1 + m_2$. Then, the amalgamation of such monomials is given by (1.2) in the form of

$$\begin{aligned} \sum_{m_1 + m_2 = n} C[m_1, m_2] t_1^{m_1} t_2^{m_2} &= [Y^n] \left(\sum_{k=1}^{3n+1} \frac{1}{k} [z^{k-1}] (1 + Y t_1 z^2 + Y t_2 z^3)^k \right) \\ &= [Y^n] \sum_{m_1, m_2 \geq 0} \frac{\binom{1+2m_1+3m_2}{m_1, m_2, 1+m_1+2m_2}}{1+2m_1+3m_2} Y^{m_1+m_2} t_1^{m_1} t_2^{m_2} \\ &= \sum_{\substack{m_1, m_2 \geq 0 \\ m_1 + m_2 = n}} \frac{\binom{1+2m_1+3m_2}{m_1, m_2, 1+m_1+2m_2}}{1+2m_1+3m_2} t_1^{m_1} t_2^{m_2} \\ &= \sum_{m_2=0}^n \frac{\binom{1+2n+m_2}{n-m_2, m_2, 1+n+m_2}}{1+2n+m_2} t_1^{n-m_2} t_2^{m_2} \\ &= \sum_{k=0}^n \frac{\binom{n}{k} \binom{2n+1+k}{n+1+k}}{2n+1+k} t_1^{n-k} t_2^k. \end{aligned}$$

For example, the following reveal both coefficients $C[m_1, m_2]$ and $G[m_1, m_2]$:

$$\begin{aligned} \sum_{m_1 + m_2 = 3} C[m_1, m_2] t_1^{m_1} t_2^{m_2} &= (t_1 + t_2)(5t_1^2 + 16t_1 t_2 + 12t_2^2), \\ \sum_{m_1 + m_2 = 4} C[m_1, m_2] t_1^{m_1} t_2^{m_2} &= (t_1 + t_2)(14t_1^3 + 70t_1^2 t_2 + 110t_1 t_2^2 + 55t_2^3). \end{aligned}$$

As a first step, we reprove that the linear term $t_2 + t_3$ divides the polynomial

$$P_n(t_1, t_2) := \sum_{k=0}^n \frac{\binom{n}{k} \binom{2n+1+k}{n+1+k}}{2n+1+k} t_1^{n-k} t_2^k.$$

This is equivalent to proving that $P_n(-t_2, t_2) = 0$, which, in turn, is equivalent to the following identity:

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k} \binom{2n+1+k}{n+1+k}}{2n+1+k} = 0.$$

To continue, we invoke the role of the WZ method. Define the functions $F(n, k) := (-1)^k \frac{\binom{n}{k} \binom{2n+1+k}{n+1+k}}{2n+1+k}$ and also $H(n, k) := -F(n, k) \cdot \frac{k(n+1+k)}{n(2n+1)}$ to verify $F(n, k) = H(n, k+1) - H(n, k)$. The rest is routine [5].

Our next step will actually find $G[m_1, m_2]$. For that we perform the division $\frac{P_n(t_1, t_2)}{t_1 + t_2}$ to obtain (algebraically) that

$$\begin{aligned} [t_1^{n-1-i} t_2^i] \left(\frac{P_n(t_1, t_2)}{t_1 + t_2} \right) &= \sum_{j=0}^i (-1)^{i-j} \frac{\binom{n}{j} \binom{2n+1+j}{n+1+j}}{2n+1+j} \\ &= (-1)^i [H(n, i+1) - H(n, 0)] \\ &= (-1)^i H(n, i+1) \\ &= \frac{1}{2n+1} \binom{n-1}{i} \binom{2n+1+i}{n+1+i} \end{aligned}$$

which leads to (an equivalent form of) the first conjecture [4] on $G[m_1, m_2]$. To wit:

Theorem 1.1. *For non-negative integers m_1 and m_2 , we have*

$$G[m_1, m_2] = \frac{1}{(2m_1 + 2m_2 + 3)(m_1 + m_2 + 1)} \frac{(2m_1 + 3m_2 + 3)!}{(m_1 + 2m_2 + 2)! m_1! m_2!}.$$

2. ON THE SECOND CONJECTURE

Now that the reader, hopefully, is getting accustomed to our proof-procedure as depicted in Section 1, let's move on to next conjecture [4, p. 399] which does generalize the one we just finished proving. For brevity, denote $\widetilde{G} = \widetilde{G}[m_a, m_{a+1}] = G[0, 0, \dots, m_a, m_{a+1}]$. Again, we revive the Lagrange Inversion (1.2). Suppose $n = m_a + m_{a+1}$. Then the total content of such

monomials is encapsulated by

$$\begin{aligned}
\sum_{m_a+m_{a+1}=n} \tilde{G} t_a^{m_2} t_{a+1}^{m_3} &= \frac{[Y^n]}{t_a + t_{a+1}} \sum_{k=1}^{(a+1)n+1} \frac{1}{k} [z^{k-1}] (1 + Y t_a z^a + Y t_{a+1} z^{a+1})^k \\
&= \frac{[Y^n]}{t_a + t_{a+1}} \sum_{m_a, m_{a+1} \geq 0} \frac{\binom{1+am_a+(a+1)m_{a+1}}{m_a, m_{a+1}, 1+(a-1)m_a+am_{a+1}} Y^{m_a+m_{a+1}} t_a^{m_a} t_{a+1}^{m_{a+1}}}{1 + am_a + (a+1)m_{a+1}} \\
&= \sum_{\substack{m_a, m_{a+1} \geq 0 \\ m_a+m_{a+1}=n}} \frac{\binom{1+am_a+(a+1)m_{a+1}}{m_a, m_{a+1}, 1+(a-1)m_a+am_{a+1}}}{1 + am_a + (a+1)m_{a+1}} \frac{t_a^{m_a} t_{a+1}^{m_{a+1}}}{t_a + t_{a+1}} \\
&= \sum_{m_{a+1}=0}^n \frac{\binom{1+an+m_{a+1}}{n-m_{a+1}, m_{a+1}, 1+(a-1)n+m_3}}{1 + an + m_{a+1}} \frac{t_a^{n-m_{a+1}} t_{a+1}^{m_{a+1}}}{t_a + t_{a+1}} \\
&= \sum_{k=0}^n \frac{\binom{n}{k} \binom{an+1+k}{(a-1)n+1+k}}{an + 1 + k} \frac{t_a^{n-k} t_{a+1}^k}{t_a + t_{a+1}}.
\end{aligned}$$

As a first step, we justify that the linear term $t_a + t_{a+1}$ divides the polynomial

$$P_n(t_a, t_{a+1}) := \sum_{k=0}^n \frac{\binom{n}{k} \binom{an+1+k}{(a-1)n+1+k}}{an + 1 + k} t_a^{n-k} t_{a+1}^k.$$

This is tantamount to $P_n(-t_{a+1}, t_{a+1}) = 0$ which is equivalent to the identity that

$$\sum_{k=0}^n (-1)^k \frac{\binom{n}{k} \binom{an+1+k}{(a-1)n+1+k}}{an + 1 + k} = 0.$$

Again, apply the Wilf-Zeilberger approach with $F(n, k) := \frac{(-1)^k \binom{n}{k} \binom{an+1+k}{(a-1)n+1+k}}{an+1+k}$ and $H(n, k) := -F(n, k) \cdot \frac{k((a-1)n+1+k)}{n(an+1)}$ to verify $F(n, k) = H(n, k+1) - H(n, k)$. The rest is trivial.

Our next step will actually determine $\tilde{G}[m_a, m_{a+1}]$. To this effect, let's divide $\frac{P_n(t_a, t_{a+1})}{t_a + t_{a+1}}$ to obtain (routinely) that

$$\begin{aligned}
[t_a^{n-1-i} t_{a+1}^i] \left(\frac{P_n(t_a, t_{a+1})}{t_a + t_{a+1}} \right) &= \sum_{j=0}^i (-1)^{i-j} \frac{\binom{n}{j} \binom{an+1+j}{(a-1)n+1+j}}{an + 1 + j} \\
&= (-1)^i [H(n, i+1) - H(n, 0)] = (-1)^i H(n, i+1) \\
&= \frac{1}{an+1} \binom{n-1}{i} \binom{an+1+i}{(a-1)n+1+i}
\end{aligned}$$

which proves the desired conjecture on $\tilde{G}[m_a, m_{a+1}]$. To wit:

Theorem 2.1. Denote $m = m_a + m_{a+1}$. For integers $m_a, m_{a+1} \geq 0$ there holds

$$\widetilde{G}[m_a, m_{a+1}] = \frac{(am_a + (a+1)(m_{a+1} + 1))!}{(a(m+1) + 1)(m+1)((a-1)m_a + a(m_{a+1} + 1))!m_a!m_{a+1}!}.$$

3. ON THE THIRD CONJECTURE

The proof of the last conjecture [4, p. 399] is a bit more complicated.

Theorem 3.1. For the $2a$ -variate case, we have

$$G[-f, f, \dots, -f, f] = \sum_n a^n f^n.$$

Proof. To begin, we make a slight alteration by writing $(-1)^i t_i$ instead of the customary plain t_i [4]. Thanks to the Lagrange Inversion (1.2), we have

$$\begin{aligned} & [Y^n] \left(\sum_{k=1}^{\infty} \frac{1}{k} [z^{k-1}] \left(1 - Y t_1 z^2 + Y t_2 z^3 - \dots - Y t_{2a-1} z^{2a} + Y t_{2a} z^{2a+1} \right)^k \right) \\ &= [Y^n] \sum_{m_1, \dots, m_{2a} \geq 0} \frac{(-1)^{m_1 + \dots + m_{2a-1}} \binom{1+2m_1+3m_2+\dots+(2a+1)m_{2a}}{m_1, m_2, \dots, m_{2a}, 1+m_1+2m_2+\dots+(2a)m_{2a}} (Y t_1)^{m_1} \dots (Y t_{2a})^{m_{2a}}}{1 + 2m_1 + 3m_2 + \dots + (2a+1)m_{2a}} \\ &= \sum_{\substack{m_1, \dots, m_{2a} \geq 0 \\ m_1 + \dots + m_{2a} = n}} \frac{(-1)^{m_1 + \dots + m_{2a-1}} \binom{1+2m_1+3m_2+\dots+(2a+1)m_{2a}}{m_1, m_2, \dots, m_{2a}, 1+m_1+2m_2+\dots+(2a)m_{2a}} t_1^{m_1} \dots t_{2a}^{m_{2a}}}{1 + 2m_1 + 3m_2 + \dots + (2a+1)m_{2a}}. \end{aligned}$$

First, consider the case $a = 1$ and refer back to Theorem 1.1 (and its proof), to gather that if $t_1 = -f$ and $t_2 = f$ then, as expected, we arrive at

$$f^{n-1} \sum_{m=0}^{n-1} \frac{(-1)^{n-1-m}}{2n+1} \binom{n-1}{m} \binom{2n+1+m}{n+1+m} = f^{n-1}$$

as justified by the *WZ-certificate* [5] given by

$$R(n, m) := \frac{m(8mn + 10n^2 + 6m + 15n + 6)}{2(2n+3)(n+1)(n-m)}.$$

Second, we go back to study the above-posed calculations when $a > 1$. To set the stage, substitute $t_1 = t_2 = \dots = t_{2a-1} = f$ while leaving out t_{2a} as an indeterminate. The outcome takes the form

$$\sum_{\substack{m_1, \dots, m_{2a} \geq 0 \\ m_1 + \dots + m_{2a} = n}} \frac{(-1)^{m_1 + m_3 + \dots + m_{2a-1}} \binom{1+2m_1+3m_2+\dots+(2a+1)m_{2a}}{m_1, m_2, \dots, m_{2a}, 1+m_1+2m_2+\dots+(2a)m_{2a}} f^{n-m_{2a}} t_{2a}^{m_{2a}}}{1 + 2m_1 + 3m_2 + \dots + (2a+1)m_{2a}}.$$

At this point, divide out the current polynomial (in t_{2a}) by the linear factor

$$-t_1 + t_2 - \dots - t_{2a-3} + t_{2a-1} - t_{2a-1} + t_{2a} = t_{2a} - f$$

and then replace t_{2a} by f . That leads to the sum

$$f^{n-1} \sum_{i=0}^{n-1} \sum_{m_{2a}=0}^i \sum_{\substack{m_1, \dots, m_{2a} \geq 0 \\ m_1 + \dots + m_{2a} = n}} \frac{(-1)^{1+m_1+m_3+\dots+m_{2a-1}} \binom{1+2m_1+3m_2+\dots+(2a+1)m_{2a}}{m_1, m_2, \dots, m_{2a}, 1+m_1+2m_2+\dots+(2a)m_{2a}}}{1+2m_1+3m_2+\dots+(2a+1)m_{2a}}.$$

Therefore, our main task that remains is to prove the identity declared by

$$\sum_{i=0}^{n-1} \sum_{\substack{m_1, \dots, m_{2a-1} \geq 0 \\ m_1 + \dots + m_{2a} = n \\ 0 \leq m_{2a} \leq i}} \frac{(-1)^{1+m_1+m_3+\dots+m_{2a-1}} \binom{1+2m_1+3m_2+\dots+(2a+1)m_{2a}}{m_1, m_2, \dots, m_{2a}, 1+m_1+2m_2+\dots+(2a)m_{2a}}}{1+2m_1+3m_2+\dots+(2a+1)m_{2a}} = a^{n-1}.$$

To put this more succinctly, introduce some notation. Let \mathcal{P} denote the set of all integer partitions λ , written as $\lambda = (\lambda_1, \lambda_2, \dots)$ or $\lambda = 1^{m_1} 2^{m_2} \dots (2a)^{m_{2a}}$. The size of λ is denoted by $|\lambda| = \lambda_1 + \lambda_2 + \dots = m_1 + 2m_2 + \dots + (2a)m_{2a}$ while we use $\ell(\lambda) = m_1 + m_2 + \dots + m_{2a}$ for the length of the partition. So, the claim stands at

$$(3.1) \quad \sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda) = n \\ \lambda_1 \leq 2a}} (-1)^{1+|\lambda|} \cdot \frac{(n - m_{2a}) \binom{n}{m_1, \dots, m_{2a}} \binom{|\lambda| + n + 1}{|\lambda| + 1}}{|\lambda| + n + 1} = a^{n-1}.$$

We find it more convenient to split up this assertion into two separate claims

$$(3.2) \quad (-1)^1 \sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda) = n \\ \lambda_1 \leq 2a}} (-1)^{|\lambda|} \binom{n}{m_1, \dots, m_{2a}} \binom{|\lambda| + n}{|\lambda| + 1} = 0,$$

$$(3.3) \quad \sum_{\substack{\mu \in \mathcal{P} \\ \ell(\mu) = n-1 \\ \mu_1 \leq 2a}} (-1)^{|\mu|} \binom{n-1}{m_1, \dots, m_{2a}} \binom{|\mu| + 2a + n}{|\mu| + 2a + 1} = a^{n-1}.$$

One arrives at (3.2) due to $\frac{n \binom{|\lambda| + n + 1}{|\lambda| + 1}}{|\lambda| + n + 1} = \binom{|\lambda| + n}{|\lambda| + 1}$ and (3.3) arises because of $m_{2a} \binom{n}{m_1, \dots, m_{2a}} \frac{(|\lambda| + n)!}{(|\lambda| + 1)! n!} = \binom{n-1}{m_1, \dots, m_{2a}-1} \binom{|\lambda| + n}{|\lambda| + 1}$ and then we reindex $m'_{2a} = m_{2a} - 1$ to convert $|\lambda| = |\mu| + 2a$ where $\ell(\mu) = n - 1$.

In fact, let's generalize (3.2) and (3.3) by introducing an extra parameter x .

Claim 1: For positive integers n, a and an indeterminate x , we have

$$\sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda) = n \\ \lambda_1 \leq 2a}} (-1)^{|\lambda|} \binom{n}{m_1, \dots, m_{2a}} \binom{|\lambda| + n + x}{n - 1} = 0.$$

Claim 2: For positive integers n, a and an indeterminate x , we have

$$\sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda)=n-1 \\ \lambda_1 \leq 2a}} (-1)^{|\lambda|} \binom{n-1}{m_1, \dots, m_{2a}} \binom{|\lambda| + n + x}{n-1} = a^{n-1}.$$

Claim 2 implies Claim 1: We apply the multinomial recurrence (assume $n = k_1 + \dots + k_r$)

$$(3.4) \quad \binom{n}{k_1, \dots, k_r} = \binom{n-1}{k_1-1, \dots, k_r} + \dots + \binom{n-1}{k_1, \dots, k_r-1}$$

followed by appropriate reindexing so that

$$\begin{aligned} & \sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda)=n \\ \lambda_1 \leq 2a}} (-1)^{|\lambda|} \binom{n}{m_1, \dots, m_{2a}} \binom{|\lambda| + n + x}{n-1} \\ &= \sum_{i=1}^{2a} \sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda)=n \\ \lambda_1 \leq 2a}} (-1)^{|\lambda|} \binom{n-1}{m_1, \dots, m_i-1, \dots, m_{2a}} \binom{|\lambda| + n + x}{n-1} \\ &= \sum_{i=1}^{2a} \sum_{\substack{\mu \in \mathcal{P} \\ \ell(\mu)=n-1 \\ \mu_1 \leq 2a}} (-1)^{|\mu|+i} \binom{n-1}{m_1, \dots, m'_i, \dots, m_{2a}} \binom{|\mu| + n + (x+i)}{n-1} \\ &= \sum_{i=1}^{2a} (-1)^i \sum_{\substack{\mu \in \mathcal{P} \\ \ell(\mu)=n-1 \\ \mu_1 \leq 2a}} (-1)^{|\mu|} \binom{n-1}{m_1, \dots, m'_i, \dots, m_{2a}} \binom{|\mu| + n + (x+i)}{n-1} \\ &= a^{n-1} \sum_{i=1}^{2a} (-1)^i \\ &= 0. \end{aligned}$$

Proof of Claim 2: Let's now utilize the multinomial theorem (1.1) and constant-term extraction. Start by noting the constant-term extraction

$$\begin{aligned} \binom{|\lambda| + n + x}{n-1} &= \binom{m_1 + 2m_2 + \dots + (2a)m_{2a} + n + x}{n-1} \\ &= \mathbf{CT}_z \left[\frac{(1+z)^{m_1+2m_2+\dots+(2a)m_{2a}+n+x}}{z^{n-1}} \right]. \end{aligned}$$

Insert this into the left-hand side of Claim 2, take \mathbf{CT}_z outside the sum, factor out the inside and reapply the multinomial theorem in reverse (1.1)

to get

$$\begin{aligned}
& \sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda)=n-1 \\ \lambda_1 \leq 2a}} (-1)^{|\lambda|} \binom{n-1}{m_1, \dots, m_{2a}} \binom{|\lambda| + n + x}{n-1} \\
&= \mathbf{CT}_z \left[\frac{(1+z)^{n+x}}{z^{n-1}} \sum \binom{n-1}{m_1, \dots, m_{2a}} (-1-z)^{m_1} (-1-z)^{2m_2} \dots (-1-z)^{(2a)m_{2a}} \right] \\
&= \mathbf{CT}_z \left[\frac{(1+z)^{n+x}}{z^{n-1}} \left\{ -(1+z)^1 + (1+z)^2 - (1+z)^3 + \dots + (1+z)^{2a} \right\}^{n-1} \right].
\end{aligned}$$

Next, follow through with the geometric series expansion to obtain

$$\begin{aligned}
& \sum_{\substack{\lambda \in \mathcal{P} \\ \ell(\lambda)=n-1 \\ \lambda_1 \leq 2a}} (-1)^{|\lambda|} \binom{n-1}{m_1, \dots, m_{2a}} \binom{|\lambda| + n + x}{n-1} \\
&= \mathbf{CT}_z \left[(-1)^{n-1} \frac{(1+z)^{2n+x-1}}{z^{n-1}} \left\{ \frac{1 - (1+z)^{2a}}{2+z} \right\}^{n-1} \right] \\
&= \mathbf{CT}_z \left[\frac{(1+z)^{2n+x-1}}{(2z)^{n-1}} \left\{ \frac{z \sum_{k=1}^{2a} \binom{2a}{k} z^{k-1}}{1 + \frac{z}{2}} \right\}^{n-1} \right] = a^{n-1}.
\end{aligned}$$

The proof is indeed complete. \square

Remark 3.2. On [4, p. 399], it is stated that “With $k-2$ leading zeros, we conjecture that $G[0, \dots, m_k]$ is a two-parameter Fuss-Catalan number.” In light of the conjectures we already proved, the current claim is rather obvious (for further discussion on the topic the reader is directed to [2]).

Remark 3.3. One can prove both Theorem 1.1 and 2.1 with the following observation. It suffice to explain this for Theorem 1.1. Since $C[m_1, m_2]$ are known from the Lagrange Inversion and because we have an explicit conjectured formula $G[m_1, m_2]$ due to [4], all that is required is to verify the relation $G[m_1 - 1, m_2] + G[m_1, m_2 - 1] = C[m_1, m_2]$. This, however, is routine. Of course, the proofs in Section 1 and 2 do not assume knowing $C[m_1, m_2]$ and $G[m_1, m_2]$ *a priori*: they are pure *derivations* from scratch.

Remark 3.4. We offer (the proof is analogous to Theorem 2.1 but omitted) the assertion that

$$G[0, \dots, 0, m_s, 0, \dots, m_t] = \frac{1}{n} \sum_{j=0}^i (-1)^{i-j} \binom{n}{j} \binom{(s+1)n + (t-s)j}{n-1},$$

where we used $m_s = n-1-i$, $m_t = i$.

Remark 3.5. We also offer (the proof is analogous to Theorem 3.1 but omitted) the assertion that for a generalized $2a$ -variate case, we have

$$\begin{aligned} & G[-c_a f, c_1 f, -c_1 f, c_2 f, -c_2 f, \dots, c_{a-1} f, -c_{a-1} f, c_a f] \\ &= \sum_n (2ac_a - c_1 - c_2 - \dots - c_a)^n f^n. \end{aligned}$$

Acknowledgment. After the completion of this work, Dean Rubine informed us that he has independently proved a couple of the conjectures with a different method [3]. The authors here together with Manuel Kauers [1] are exploring the structure of the Geode in its generality.

REFERENCES

- [1] T. Amdeberhan, M. Kauers, D. Zeilberger, *in preparation*.
- [2] S. R. Mane, *Multiparameter Fuss–Catalan numbers with application to algebraic equations*, <https://arxiv.org/pdf/1607.04144>
- [3] D. Rubine, *private communication*
- [4] N. J. Wildberger, D. Rubine, *A Hyper-Catalan Series Solution to Polynomial Equations, and the Geode*, Amer. Math. Monthly, **132**, (2025), no. 5, 383-402. <https://www.tandfonline.com/doi/full/10.1080/00029890.2025.2460966>
- [5] H. S. Wilf, D. Zeilberger, *Rational functions certify combinatorial identities*, J. Amer. Math. Soc. **3** (1990), no. 1, 147-158. <https://www.ams.org/journals/jams/1990-03-01/S0894-0347-1990-1007910-7/S0894-0347-1990-1007910-7.pdf>
- [6] D. Zeilberger, *Lagrange Inversion Without Tears (Analysis) (based on Henrici)*, <https://sites.math.rutgers.edu/~zeilberg/mamarim/mamarimPDF/lag.pdf>

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118, USA
Email address: tamdeber@tulane.edu

RUTGERS UNIVERSITY, DEPARTMENT OF MATHEMATICS, 110 FRELINGHUYSEN RD, PISCATAWAY, NJ 08854, USA
Email address: DoronZeil@gmail.com