

IDENTITIES FOR GENERALIZED EULER POLYNOMIALS

LIN JIU, VICTOR H. MOLL, AND CHRISTOPHE VIGNAT

ABSTRACT. For $N \in \mathbb{N}$, let T_N be the Chebyshev polynomial of the first kind. Expressions for the sequence of numbers $p_\ell^{(N)}$, defined as the coefficients in the expansion of $1/T_N(1/z)$, are provided. These coefficients give formulas for the classical Euler polynomials in terms of the so-called generalized Euler polynomials. The proofs are based on a probabilistic interpretation of the generalized Euler polynomials recently given by Klebanov et al. Asymptotics of $p_\ell^{(N)}$ are also provided.

1. INTRODUCTION

The Euler numbers E_n , defined by the generating function

$$(1.1) \quad \frac{1}{\cosh z} = \sum_{n=0}^{\infty} E_n \frac{z^n}{n!}$$

and the Euler polynomials $E_n(x)$ that generalize them

$$(1.2) \quad \sum_{n=0}^{\infty} E_n(x) \frac{z^n}{n!} = \frac{2e^{xz}}{e^z + 1}$$

([2, 9.630,9.651]) are examples of basic special functions. It follows directly from the definition that $E_n = 0$ for n odd. Moreover, the relation $E_n = 2^n E_n(\frac{1}{2})$ follows by setting $x = \frac{1}{2}$ in (1.2), replacing z by $2z$ and comparing with (1.1).

Moreover, the identity

$$(1.3) \quad \frac{2e^{xz}}{e^z + 1} = \frac{2e^{(x-1/2)z}}{e^{z/2} + e^{-z/2}}$$

produces

$$(1.4) \quad E_n(x) = \sum_{k=0}^n \binom{n}{k} \frac{E_k}{2^k} \left(x - \frac{1}{2}\right)^{n-k} = \sum_{k=0}^n \binom{n}{k} E_k \left(\frac{1}{2}\right) \left(x - \frac{1}{2}\right)^{n-k},$$

that gives $E_n(x)$ in terms of the Euler numbers (see [2, 9.650]).

Date: January 30, 2014.

2010 Mathematics Subject Classification. Primary 11B68, Secondary 60E05.

Key words and phrases. Generalized Euler polynomials, hyperbolic secant distributions, Chebyshev polynomials.

The *generalized Euler polynomials* $E_n^{(p)}(z)$, defined by the generating function

$$(1.5) \quad \sum_{n=0}^{\infty} E_n^{(p)}(x) \frac{z^n}{n!} = \left(\frac{2}{1+ez} \right)^p e^{xz}, \quad \text{for } p \in \mathbb{N}$$

are polynomials extending $E_n(x)$, the case $p = 1$. These appear in Section 24.16 of [5]. The definition leads directly to the expression

$$(1.6) \quad E_n^{(p)}(x) = \sum_{k=0}^n \binom{n}{k} x^k E_{n-k}^{(p)}(0),$$

where the *generalized Euler numbers* $E_n^{(p)}(0)$ are defined recursively by

$$(1.7) \quad E_n^{(p)}(0) = \sum_{k=0}^n \binom{n}{k} E_k^{(p-1)}(0) E_{n-k}(0),$$

for $p > 1$ and initial condition $E_n^{(1)}(0) = E_n(0)$.

2. A PROBABILISTIC REPRESENTATION OF EULER POLYNOMIALS AND THEIR GENERALIZATIONS

This section discusses probabilistic representations of the Euler polynomials and their generalizations. The results involve the expectation operator \mathbb{E} defined by

$$(2.1) \quad \mathbb{E}g(L) = \int g(x) f_L(x) dx,$$

with $f_L(x)$ the probability density of the random variable L and for any function g such that the integral exists.

Proposition 2.1. *Let L be a random variable with hyperbolic secant density*

$$(2.2) \quad f_L(x) = \operatorname{sech} \pi x, \quad \text{for } x \in \mathbb{R}.$$

Then the Euler polynomial is given by

$$(2.3) \quad E_n(x) = \mathbb{E} \left(x + \imath L - \frac{1}{2} \right)^n.$$

Proof. The right hand-side of (2.3) is

$$\begin{aligned} \mathbb{E} \left(x + \imath L - \frac{1}{2} \right)^n &= \int_{-\infty}^{\infty} \left(x - \frac{1}{2} + \imath t \right)^n \operatorname{sech} \pi t dt \\ &= \sum_{j=0}^n \binom{n}{j} \left(x - \frac{1}{2} \right)^{n-j} \imath^j \int_{-\infty}^{\infty} t^j \operatorname{sech} \pi t dt \end{aligned}$$

The identity

$$(2.4) \quad \int_{-\infty}^{\infty} t^k \operatorname{sech} \pi t dt = \frac{|E_k|}{2^k}$$

holds for k odd, since both sides vanish and for k even, it appears as entry 3.523.4 in [2]. A proof of this entry may be found in [1]. Then, using $|E_{2n}| = (-1)^n E_{2n}$ (entry 9.633 in [2])

$$(2.5) \quad \mathbb{E}(x + \iota L - \frac{1}{2})^n = \sum_{j=0}^n \binom{n}{j} (x - \frac{1}{2})^{n-j} \frac{E_j}{2^j} = E_n(x).$$

□

There is a natural extension to the case of $E_n^{(p)}(x)$. The proof is similar to the previous case, so it is omitted.

Theorem 2.2. *Let $p \in \mathbb{N}$ and L_j , $1 \leq j \leq p$ a collection of independent identically distributed random variables with hyperbolic secant distribution. Then*

$$(2.6) \quad E_n^{(p)}(x) = \mathbb{E} \left[x + \sum_{j=1}^p (\iota L_j - \frac{1}{2}) \right]^n.$$

In a recent paper, L. B. Klebanov et al. [3] considered random sums of independent random variables of the form

$$(2.7) \quad \frac{1}{N} \sum_{j=1}^{\mu_N} L_j$$

where the random number of summands μ_N is independent of the L_j 's and is described below.

Definition 2.3. Let $N \in \mathbb{N}$ and $T_N(z)$ be the Chebyshev polynomial of the first kind. The random variable μ_N taking values in \mathbb{N} , is defined by its generating function

$$(2.8) \quad \mathbb{E}z^{\mu_N} = \frac{1}{T_N(1/z)}.$$

Information about the Chebyshev polynomials appears in [2] and [5].

Example 2.4. Take $N = 2$. Then $T_2(z) = 2z^2 - 1$ gives

$$(2.9) \quad \mathbb{E}z^{\mu_2} = \frac{1}{T_2(1/z)} = \frac{z^2}{2 - z^2} = \sum_{\ell=1}^{\infty} \frac{z^{2\ell}}{2^\ell}.$$

Therefore μ_2 takes the value 2ℓ , with $\ell \in \mathbb{N}$, with probability

$$(2.10) \quad \Pr(\mu_2 = 2\ell) = 2^{-\ell}.$$

In [3], Klebanov et al. prove the following result.

Theorem 2.5 (Klebanov et al.). *Assume $\{L_j\}$ is a sequence of independent identically distributed random variables with hyperbolic secant distribution. Then, for all $N \geq 2$ and μ_N defined in (2.8), the random variable*

$$(2.11) \quad L := \frac{1}{N} \sum_{j=1}^{\mu_N} L_j$$

has the same hyperbolic secant distribution.

3. THE EULER POLYNOMIALS IN TERMS OF THE GENERALIZED ONES

The identities (1.6) and (1.7) can be used to express the generalized Euler polynomial $E_n^{(p)}(x)$ in terms of the standard Euler polynomials $E_n(x)$. However, to the best of our knowledge, there is no formula that allows to express $E_n(x)$ in terms of $E_n^{(p)}(x)$. This section presents such a formula.

Definition 3.1. Let $N \in \mathbb{N}$. The sequence $\{p_\ell^{(N)} : \ell = 0, 1, \dots\}$ is defined as the coefficients in the expansion

$$(3.1) \quad \frac{1}{T_N(1/z)} = \sum_{\ell=0}^{\infty} p_\ell^{(N)} z^\ell.$$

Definition 2.3 shows that

$$(3.2) \quad p_\ell^{(N)} = \Pr(\mu_N = \ell), \quad \text{for } \ell \in \mathbb{N}.$$

The numbers $p_\ell^{(N)}$ will be referred as the *probability numbers*.

Example 3.2. For $N = 2$, Example 2.4 gives

$$(3.3) \quad p_\ell^{(2)} = \begin{cases} 0 & \text{if } \ell \text{ is odd} \\ 2^{-\ell/2} & \text{if } \ell \text{ is even, } \ell \neq 0. \end{cases}$$

The coefficients $p_\ell^{(N)}$ are now used to produce expansions of $E_n(x)$, one for each $N \in \mathbb{N}$, in terms of the generalized Euler polynomials.

Theorem 3.3. *The Euler polynomials satisfy, for all $N \in \mathbb{N}$,*

$$(3.4) \quad E_n(x) = \frac{1}{N^n} \mathbb{E} \left[E_n^{(\mu_N)} \left(\frac{1}{2} \mu_N + N \left(x - \frac{1}{2} \right) \right) \right]$$

Proof. From (2.3) and (2.11)

$$(3.5) \quad E_n \left(\frac{1}{2} \right) = \mathbb{E}(\imath L)^n = \frac{1}{N^n} \mathbb{E} \left[\imath \sum_{j=1}^{\mu_N} L_j \right]^n,$$

with Theorem 2.2, this yields

$$(3.6) \quad \mathbb{E} \left[E_n^{(\mu_N)} \left(\frac{\mu_N}{2} \right) \right] = \mathbb{E} \left[\imath \sum_{j=1}^{\mu_N} L_j \right]^n = N^n E_n \left(\frac{1}{2} \right).$$

Using identity (1.4), it follows that

$$\begin{aligned}
E_n(x) &= \sum_{k=0}^n \binom{n}{k} E_k\left(\frac{1}{2}\right) \left(x - \frac{1}{2}\right)^{n-k} \\
&= \mathbb{E} \left[\sum_{k=0}^n \binom{n}{k} N^{-k} E_k^{(\mu_N)}\left(\frac{1}{2}\mu_N\right) \left(x - \frac{1}{2}\right)^{n-k} \right] \\
&= \mathbb{E} \left[\sum_{k=0}^n \binom{n}{k} N^{-k} (\imath L_1 + \cdots + \imath L_{\mu_N})^k \left(x - \frac{1}{2}\right)^{n-k} \right] \\
&= \mathbb{E} \left[\frac{1}{N^n} \sum_{k=0}^n \binom{n}{k} (\imath L_1 + \cdots + \imath L_{\mu_N})^k \left(N\left(x - \frac{1}{2}\right)\right)^{n-k} \right] \\
&= \mathbb{E} \left[\frac{1}{N^n} (\imath L_1 + \cdots + \imath L_{\mu_N} + N\left(x - \frac{1}{2}\right))^n \right] \\
&= \mathbb{E} \left[\frac{1}{N^n} (\imath L_1 + \cdots + \imath L_{\mu_N} + z - \frac{1}{2}\mu_N)^n \right] \\
&= \frac{1}{N^n} \mathbb{E} \left[E_n^{(\mu_N)}(z) \right],
\end{aligned}$$

where $z = \frac{1}{2}\mu_N + N\left(x - \frac{1}{2}\right)$. This completes the proof. \square

The next result is established using the fact that the expectation operator \mathbb{E} satisfies

$$(3.7) \quad \mathbb{E}[h(\mu_N)] = \sum_{k=0}^{\infty} p_k^{(N)} h(k),$$

for any function h such that the right-hand side exists.

Corollary 3.4. *The Euler polynomials satisfy*

$$(3.8) \quad E_n(x) = \frac{1}{N^n} \sum_{k=N}^{\infty} p_k^{(N)} E_n^{(k)}\left(\frac{1}{2}k + N\left(x - \frac{1}{2}\right)\right).$$

Note 3.5. Corollary 3.4 gives an infinite family of expressions for $E_n(x)$ in terms of the generalized Euler polynomials $E_n^{(k)}(x)$, one for each value of $N \geq 2$.

Example 3.6. The expansion (3.8) with $N = 2$ gives

$$(3.9) \quad E_n(x) = \frac{1}{2^n} \sum_{\ell=1}^{\infty} \frac{1}{2^\ell} E_n^{(2^\ell)}(\ell + 2x - 1).$$

For instance, when $n = 1$,

$$(3.10) \quad E_1(x) = \frac{1}{2} \sum_{\ell=1}^{\infty} \frac{1}{2^\ell} E_1^{(2^\ell)}(\ell + 2x - 1)$$

and the value $E_1^{(\ell)}(x) = x - \frac{\ell}{2}$ gives

$$(3.11) \quad E_1(x) = \frac{1}{2} \sum_{\ell=1}^{\infty} \frac{1}{2^{\ell}} (\ell + 2x - 1 - \ell) = x - \frac{1}{2}$$

as expected.

4. THE PROBABILITY NUMBERS

For fixed $N \in \mathbb{N}$, the random variable μ_N has been defined by its moment generating function

$$(4.1) \quad \mathbb{E}z^{\mu_N} = \frac{1}{T_N(1/z)} = \sum_{\ell=0}^{\infty} p_{\ell}^{(N)} z^{\ell}.$$

This section presents properties of the probability numbers $p_{\ell}^{(N)}$ that appear in Corollary 3.4.

For small N , the coefficients $p_{\ell}^{(N)}$ can be computed directly by expanding the rational function $1/T_N(1/z)$ in partial fractions. Example 2.4 gave the case $N = 2$. The cases $N = 3$ and $N = 4$ are presented below.

Example 4.1. For $N = 3$, the Chebyshev polynomial is

$$(4.2) \quad T_3(z) = 4z^3 - 3z = 4z(z - \alpha)(z + \alpha),$$

with $\alpha = \sqrt{3}/2$. This yields

$$(4.3) \quad \frac{1}{T_3(1/z)} = \frac{z^3}{4(1 - \alpha z)(1 + \alpha z)} = \sum_{k=0}^{\infty} \frac{3^k}{2^{2k+2}} z^{2k+3}.$$

It follows that $p_{\ell}^{(3)} = 0$ unless $\ell = 2k + 3$ and

$$(4.4) \quad p_{2k+3}^{(3)} = \frac{3^k}{2^{2k+2}}.$$

Corollary 3.4 now gives

$$(4.5) \quad E_n(x) = \frac{1}{3^n} \sum_{k=0}^{\infty} \frac{3^k}{2^{2k+2}} E_n^{(2k+3)}(3x + k),$$

a companion to (3.9).

Example 4.2. The probability numbers for $N = 4$ are computed from the expression

$$(4.6) \quad \frac{1}{T_4(1/z)} = \frac{z^4}{z^4 - 8z^2 + 8}.$$

The factorization

$$(4.7) \quad z^4 - 8z^2 + 8 = (z^2 - \beta)(z^2 - \gamma)$$

with $\beta = 2(2 + \sqrt{2})$ and $\gamma = 2(2 - \sqrt{2})$ and the partial fraction decomposition

$$(4.8) \quad \frac{z^4}{z^4 - 8z^2 + 8} = \frac{\beta}{\beta - \gamma} \frac{1}{1 - \beta/z^2} - \frac{\gamma}{\beta - \gamma} \frac{1}{1 - \gamma/z^2}$$

show that $p_\ell^{(4)} = 0$ for ℓ odd or $\ell = 2$ and

$$(4.9) \quad p_{2\ell}^{(4)} = \frac{\sqrt{2}}{2^{2\ell+1}} \left[(2 + \sqrt{2})^{\ell-1} - (2 - \sqrt{2})^{\ell-1} \right]$$

for $\ell \geq 2$. Corollary 3.4 now gives

$$(4.10) \quad E_n(x) = \sqrt{2} \sum_{\ell=2}^{\infty} \frac{[(2 + \sqrt{2})^{\ell-1} - (2 - \sqrt{2})^{\ell-1}]}{2^{2\ell+1}} E_n^{(2\ell)}(4x + \ell - 2).$$

Some elementary properties of the probability numbers are presented next.

Proposition 4.3. *The probability numbers $p_\ell^{(N)}$ vanish if $\ell < N$.*

Proof. The Chebyshev polynomial $T_N(z)$ has the form $2^{N-1}z^N +$ lower order terms. Then the expansion of $1/T_N(1/z)$ has a zero of order N at $z = 0$. This proves the statement. \square

Proposition 4.4. *The probability numbers $p_\ell^{(N)}$ vanish if $\ell \not\equiv N \pmod{2}$.*

Proof. The polynomial $T_N(z)$ has the same parity as N . The same holds for the rational function $1/T_N(1/z)$. \square

An expression for the probability numbers is given next.

Theorem 4.5. *Let $N \in \mathbb{N}$ be fixed and define*

$$(4.11) \quad \theta_k^{(N)} = \frac{(2k-1)\pi}{2N}.$$

Then

$$(4.12) \quad p_\ell^{(N)} = \frac{1}{N} \sum_{k=1}^N (-1)^{k+1} \sin \theta_k^{(N)} \cos^{\ell-1} \theta_k^{(N)}.$$

Proof. The Chebyshev polynomial is defined by $T_N(\cos \theta) = \cos(N\theta)$, so its roots are $z_k^{(N)} = \cos \theta_k^{(N)}$, with $\theta_k^{(N)}$ as above. The leading coefficient of $T_N(z)$ is 2^{N-1} , thus

$$(4.13) \quad \frac{1}{T_N(z)} = \frac{2^{1-N}}{\prod_{k=1}^N (z - z_k)}.$$

In the remainder of the proof, the superscript N has been dropped from $z_k^{(N)}$ and $\theta_k^{(N)}$, for clarity. Define

$$(4.14) \quad Q(z) = \prod_{k=1}^N (z - z_k).$$

The roots z_k of Q are distinct, therefore

$$(4.15) \quad \frac{1}{Q(z)} = \sum_{k=1}^N \frac{1}{Q'(z_k)} \frac{1}{z - z_k}.$$

The identity $T'_N(z) = NU_{N-1}(z)$ gives

$$(4.16) \quad Q'(z_k) = N2^{1-N}U_{N-1}(z_k)$$

where $U_j(z)$ is the Chebyshev polynomial of the second kind defined by

$$(4.17) \quad U_N(\cos \theta) = \frac{\sin(N+1)\theta}{\sin \theta}.$$

Then

$$(4.18) \quad U_{N-1}(z_k) = U_{N-1}(\cos \theta_k) = \frac{\sin N\theta_k}{\sin \theta_k}.$$

and the value $\sin N\theta_k = (-1)^{k+1}$ yields

$$(4.19) \quad Q'(z_k) = \frac{(-1)^{k+1}}{\sin \theta_k} N2^{1-N}.$$

Therefore (4.15) now gives

$$(4.20) \quad \frac{1}{Q(z)} = \frac{2^{N-1}}{N} \sum_{k=1}^N \frac{(-1)^{k+1} \sin \theta_k}{z - \cos \theta_k}.$$

It follows that

$$\begin{aligned} \frac{1}{T_N(1/z)} &= \frac{2^{1-N}}{Q(1/z)} = \frac{1}{N} \sum_{k=1}^N (-1)^{k+1} \frac{z \sin \theta_k}{1 - z \cos \theta_k} \\ &= \frac{1}{N} \sum_{k=1}^N (-1)^{k+1} \sin \theta_k \sum_{\ell=0}^{\infty} z^{\ell+1} \cos^\ell \theta_k \\ &= \frac{1}{N} \sum_{\ell=0}^{\infty} z^{\ell+1} \sum_{k=1}^N (-1)^{k+1} \sin \theta_k \cos^\ell \theta_k. \end{aligned}$$

The proof is complete. \square

The next result provides another explicit formula for the probability numbers. The coefficients $A(n, k)$ appear in OEIS entry A008315, as entries of the Catalan triangle.

Theorem 4.6. *Let $A(n, k) = \binom{n}{k} - \binom{n}{k-1}$. Then, if $N \equiv \ell \pmod{2}$,*

$$p_\ell^{(N)} = \frac{1}{2^\ell} \sum_{t=\lfloor \frac{1}{2}(\frac{2-\ell}{N}-1) \rfloor}^{\lfloor \frac{1}{2}(\frac{\ell}{N}-1) \rfloor} (-1)^t A(\ell-1, \frac{1}{2}(\ell-(2t+1)N)),$$

indent when ℓ is not an odd multiple of N and

$$p_\ell^{(N)} = \frac{1}{2^\ell} \left[\sum_{s=1}^{\lfloor \ell/N-1 \rfloor} (-1)^{k-s} A(\ell-1, sN) \right] + \frac{(-1)^k}{2^{\ell-1}}, \text{ with } k = \frac{1}{2}(\ell/N - 1)$$

otherwise.

The proof begins with a preliminary result.

Lemma 4.7. *Let $N \in \mathbb{N}$ and $\theta_k = \frac{\pi}{2} \frac{(2k-1)}{N}$. Then*

$$(4.21) \quad f_N(z) = \sum_{k=1}^N (-1)^{k+1} e^{i\theta_k z}$$

is given by

$$(4.22) \quad f_N(z) = \frac{1 - (-1)^N e^{\pi i z}}{2 \cos\left(\frac{\pi z}{2N}\right)} \quad \text{if } z \neq (2t+1)N \text{ with } t \in \mathbb{Z}$$

and

$$(4.23) \quad f_N(z) = (-1)^t N i \quad \text{if } z = (2t+1)N \text{ for some } t \in \mathbb{Z}.$$

In particular

$$(4.24) \quad f_N(k) = \begin{cases} (-1)^{(k/N-1)/2} N i & \text{if } \frac{k}{N} \text{ is an odd integer} \\ \frac{1 - (-1)^{N+k}}{2 \cos\left(\frac{\pi k}{2N}\right)} & \text{otherwise.} \end{cases}$$

Proof. The function f_N is the sum of a geometric progression. The formula (4.23) comes from (4.22) by passing to the limit. \square

The proof of Theorem 4.6 is given now.

Proof. The expression for $p_\ell^{(N)}$ given in Theorem 4.5 yields

$$\begin{aligned} p_\ell^{(N)} &= \frac{1}{N} \sum_{k=1}^N (-1)^{k+1} \frac{(e^{i\theta_k} - e^{-i\theta_k})}{2i} \left(\frac{e^{i\theta_k} + e^{-i\theta_k}}{2} \right)^{\ell-1} \\ &= \frac{1}{2^\ell N i} \sum_{k=1}^N (-1)^{k+1} \sum_{r=0}^{\ell-1} \binom{\ell-1}{r} \left[e^{i(\ell-2r)\theta_k} - e^{i(\ell-2r-2)\theta_k} \right] \\ &= \frac{1}{2^\ell N i} \sum_{r=0}^{\ell-1} \binom{\ell-1}{r} [f_N(\ell-2r) - f_N(\ell-2r-2)] \\ &= \frac{1}{2^\ell N i} \left[\sum_{r=1}^{\ell-1} A(\ell-1, r) f_N(\ell-2r) + f_N(\ell) - f_N(-\ell) \right]. \end{aligned}$$

Now $f_N(\ell) = f_N(-\ell) = 0$ if ℓ/N is not an odd integer. On the other hand, if $\ell = (2t+1)N$, with $t \in \mathbb{Z}$, then

$$(4.25) \quad f_N(\ell) = (-1)^t N i \text{ and } f_N(-\ell) = -(-1)^t N i.$$

Thus

$$f_N(\ell) - f_N(-\ell) = \begin{cases} 2N\iota(-1)^{(\ell/N-1)/2} & \text{if } \ell \text{ is an odd multiple of } N \\ 0 & \text{otherwise.} \end{cases}$$

The simplification of the previous expression for $p_\ell^{(N)}$ is divided in two cases, according to whether ℓ is an odd multiple of N or not.

Case 1. Assume ℓ is not an odd multiple of N . Then

$$(4.26) \quad p_\ell^{(N)} = \frac{1}{2^\ell N \iota} \sum_{r=0}^{\ell-1} A(\ell-1, r) f_N(\ell-2r).$$

Moreover,

$$(4.27) \quad f_N(\ell-2r) = \begin{cases} (-1)^t N \iota & \text{if } \frac{\ell-2r}{N} = 2t+1 \\ 0 & \text{otherwise.} \end{cases}$$

Therefore

$$(4.28) \quad p_\ell^{(N)} = \frac{1}{2^\ell} \sum_{\substack{t=\frac{1}{2}(\frac{\ell}{N}-1) \\ \ell-2r=(2t+1)N}}^{\frac{1}{2}(\frac{\ell}{N}-1)} (-1)^t A(\ell-1, r).$$

Observe that $\ell - (2t+1)N$ is always an even integer, thus the index r may be eliminated from the previous expression to obtain

$$(4.29) \quad p_\ell^{(N)} = \frac{1}{2^\ell} \sum_{t=\lfloor \frac{1}{2}(\frac{\ell}{N}-1) \rfloor}^{\lfloor \frac{1}{2}(\frac{\ell}{N}-1) \rfloor} (-1)^t A(\ell-1, \frac{1}{2}(\ell - (2t+1)N)).$$

Case 2. Assume ℓ is an odd multiple of N , say $\ell = (2k+1)N$. Then

$$\begin{aligned} p_\ell^{(N)} &= \frac{1}{2^\ell N \iota} \left[\sum_{r=0}^{\ell-1} A(\ell-1, r) f_N(\ell-2r) + 2N\iota(-1)^k \right] \\ &= \frac{1}{2^\ell N \iota} \left[\sum_{r=0}^{\ell-1} A(\ell-1, r) f_N(\ell-2r) \right] + \frac{(-1)^k}{2^{\ell-1}}. \end{aligned}$$

The term $f_N(\ell-2r)$ vanishes unless $\ell-2r$ is an odd multiple of N . Given that $\ell = (2k+1)N$, the term is non-zero provided $2r$ is an even multiple of N ; say $r = sN$ for $s \in \mathbb{N}$. The range of s is $1 \leq s \leq \frac{\ell-1}{N} = 2k+1 - \frac{1}{N}$. This implies $1 \leq s \leq 2k = \ell/N - 1$, and it follows that

$$p_\ell^{(N)} = \frac{1}{2^\ell} \left[\sum_{s=1}^{\ell/N-1} (-1)^{k-s} A(\ell-1, sN) \right] + \frac{(-1)^k}{2^{\ell-1}}, \text{ with } k = \frac{1}{2}(\ell/N - 1).$$

The proof is complete. \square

Note 4.8. The expression in Theorem 4.6 shows that $p_\ell^{(N)}$ is a rational number with a denominator a power of 2 of exponent at most ℓ . Arithmetic properties of these coefficients will be described in a future publication [4]. Moreover, the probability numbers $p_\ell^{(N)}$ appear in the description of a random walk on N sites. Details will appear in [4].

5. AN ASYMPTOTIC EXPANSION

The final result deals with the asymptotic behavior of the probability numbers $p_\ell^{(N)}$.

Theorem 5.1. *Let $\varphi_N(z) = \mathbb{E}[z^{\mu_N}]$. Then, for fixed z in the unit disk $|z| < 1$,*

$$(5.1) \quad \varphi_N(z) \sim \left(\frac{z}{1 + \sqrt{1 - z^2}} \right)^N, \text{ as } N \rightarrow \infty.$$

Proof. The generating function satisfies

$$(5.2) \quad \varphi_N(z) = 1/T_N(1/z) = \frac{z^N}{2^{N-1}} \prod_{k=1}^N \left(1 - z \cos \theta_k^{(N)} \right)^{-1}$$

with $\theta_k^{(N)} = (2k - 1)\pi/2N$ as before. Then

$$(5.3) \quad \log \varphi_N(z) = \log 2 + N \log \frac{z}{2} - \sum_{k=1}^N \log \left(1 - z \cos \theta_k^{(N)} \right).$$

The last sum is approximated by a Riemann integral

$$\frac{1}{N} \sum_{k=1}^N \log \left(1 - z \cos \theta_k^{(N)} \right) \sim \frac{1}{\pi} \int_0^\pi \log(1 - z \cos \theta) d\theta = \log \left(\frac{1 + \sqrt{1 - z^2}}{2} \right).$$

The last evaluation is elementary. It appears as entry 4.224.9 in [2]. It follows that

$$(5.4) \quad \log \varphi_N(z) \sim \log 2 + N \log \left(\frac{z}{2} \right) - N \log \left(\frac{1 + \sqrt{1 - z^2}}{2} \right)$$

indent and this is equivalent to the result. □

The function

$$(5.5) \quad A(z) = \frac{2}{1 + \sqrt{1 - 4z}} = \sum_{n=0}^{\infty} C_n z^n$$

is the generating function for the Catalan numbers

$$(5.6) \quad C_n = \frac{1}{n+1} \binom{2n}{n}.$$

The final result follows directly from the expansion of Binet's formula for Chebyshev polynomial

$$(5.7) \quad T_N(z) = \frac{(z - \sqrt{z^2 - 1})^N + (z + \sqrt{z^2 - 1})^N}{2}.$$

Some standard notation is recalled. Given two sequences $\mathbf{a} = \{a_n\}$, $\mathbf{b} = \{b_n\}$, their convolution $\mathbf{c} = \mathbf{a} * \mathbf{b}$ is the sequence $\mathbf{c} = \{c_n\}$, with

$$(5.8) \quad c_n = \sum_{j=0}^n a_j b_{n-j}.$$

The *convolution power* $\mathbf{c}^{(*N)}$ is the convolution of \mathbf{c} with itself, N times.

Theorem 5.2. *For $N \in \mathbb{N}$ fixed, the first N nonzero terms of the sequence $q_\ell^{(N)} = 2^{\ell-1} p_\ell^{(N)}$ agree with the first N terms of the N -th convolution power $C_n^{(*N)}$ of the Catalan sequence:*

$$q_N^{(N)} = C_0^{(*N)}, q_{N+2}^{(N)} = C_1^{(*N)}, \dots, q_{N+2k}^{(N)} = C_k^{(*N)}, \dots, q_{3N-2}^{(N)} = C_{N-1}^{(*N)}.$$

In terms of generating functions, this is equivalent to

$$(5.9) \quad \left(\sum_{n=0}^{\infty} C_n z^{2n+1} \right)^N - \sum_{\ell=0}^{\infty} q_\ell^{(N)} z^\ell \sim 2^N z^{3N}.$$

Acknowledgments. The second author acknowledges the partial support of NSF-DMS 1112656. The first author is a graduate student, funded in part by the same grant.

REFERENCES

- [1] K. Boyadzhiev and V. Moll. The integrals in Gradshteyn and Ryzhik. Part 21: Hyperbolic functions. *Scientia*, 22:109–127, 2013.
- [2] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products*. Edited by A. Jeffrey and D. Zwillinger. Academic Press, New York, 7th edition, 2007.
- [3] L. B. Klebanov, A. V. Kakosyan, S. T. Rachev, and G. Temnov. On a class of distributions stable under random summation. *J. Appl. Prob.*, 49:303–318, 2012.
- [4] V. Moll and C. Vignat. Arithmetic properties of a probability sequence. *In preparation*, 2014.
- [5] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST Handbook of Mathematical Functions*. Cambridge University Press, 2010.

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118
E-mail address: ljiu@tulane.edu

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, NEW ORLEANS, LA 70118
E-mail address: vhm@tulane.edu

LSS-SUPELEC, UNIVERSITÉ ORSAY PARIS SUD 11, FRANCE
E-mail address: cvignat@tulane.edu
E-mail address: christophe.vignat@u-psud.fr